

UNITED STATES PATENT APPLICATION

PHARMACEUTICAL COMPOSITION AND METHOD FOR  
TREATING HYPOGONADISM

Inventors:

Robert E. Dudley


Olivier Palatchi

S. George Kottayil

CERTIFICATE OF MAILING BY "EXPRESS MAIL"  
UNDER 37 C.F.R. § 1.10

"Express Mail" mailing label number: EL691889850US  
Date of Mailing: October 19, 2001

I hereby certify that this correspondence is being deposited with the United States Postal Service, utilizing the "Express Mail Post Office to Addressee" service addressed to Box PATENT APPLICATION, Assistant Commissioner for Patents, Washington, DC 20231 and mailed on the above Date of Mailing with the above "Express Mail" mailing label number.

  
\_\_\_\_\_  
~~Timothy Hubalik~~ Joseph A. Mahoney

Signature Date: October 19, 2001

**PHARMACEUTICAL COMPOSITION AND METHOD FOR  
TREATING HYPOGONADISM**

**FIELD OF THE INVENTION**

5       The present invention is directed to a pharmaceutical composition comprising testosterone in a gel formulation, and to methods of using the same.

14111796V-2

## BACKGROUND OF THE INVENTION

### A. Testosterone Metabolism in Men

Testosterone is the major circulating androgen in men. More than 95% of the 6-7 mg of testosterone produced per day is secreted by the approximately 500 million Leydig cells in the testes. Two hormones produced by the pituitary gland, luteinizing hormone ("LH") and follicle stimulating hormone ("FSH"), are required for the development and maintenance of testicular function.

The most important hormone for the regulation of Leydig cell number and function is LH. In eugonadal men, LH secretion from the pituitary is inhibited through a negative-feedback pathway by increased concentrations of testosterone through the inhibition of the release of gonadotropin-releasing hormone ("GRH") by the hypothalamus. FSH promotes spermatogenesis and is essential for the normal maturation of sperm. FSH secretion from the pituitary normally is inhibited through a negative-feedback pathway by increased testosterone concentrations.

Testosterone is responsible primarily for the development and maintenance of secondary sex characteristics in men. In the body, circulating testosterone is metabolized to various 17-keto steroids through two different pathways. Testosterone can be metabolized to dihydrotestosterone ("DHT") by the enzyme  $5\alpha$ -reductase. There are two forms of  $5\alpha$ -reductase in the body: one form is found predominately in the liver and non-genital skin while another form is found in the urogenital tract of the male and the genital skin of both sexes. Testosterone can also be metabolized to estradiol ("E<sub>2</sub>") by an aromatase enzyme complex found in the liver, fat, and the testes.

Testosterone circulates in the blood 98% bound to protein. In men, approximately 40% of the binding is to the high-affinity sex hormone binding globulin ("SHBG"). The remaining

60% is bound weakly to albumin. Thus, a number of measurements for testosterone are available from clinical laboratories. The term “free” testosterone as used herein refers to the fraction of testosterone in the blood that is not bound to protein. The term “total testosterone” or “testosterone” as used herein means the free testosterone plus protein-bound testosterone. The term “bioavailable testosterone” as used herein refers to the non-SHBG bound testosterone and includes that weakly bound to albumin.

The conversion of testosterone to DHT is important in many respects. For example, DHT binds with greater affinity to SHBG than does testosterone. In addition, in many tissues, the activity of testosterone depends on the reduction to DHT, which binds to cytosol receptor proteins. The steroid-receptor complex is then transported to the nucleus where it initiates transcription and cellular changes related to androgen action. DHT is also thought to lower prostate volume and inhibit tumor development in the prostate. Thus, given the importance of DHT and testosterone in normal body functioning, researchers frequently assess and report androgen concentrations in patients as total androgen (“DHT +T”) or as a ratio of DHT to testosterone (“DHT/T ratio”).

The following table from the UCLA-Harbor Medical Center summarizes the hormone concentrations in normal adult men range:

**Table 1: Hormone Levels in Normal Men**

Hormone	Normal Range
Testosterone	298 to 1043 ng/dL
Free testosterone	3.5 to 17.9 ng/dL
DHT	31 to 193 ng/dL
DHT/T Ratio	0.052 to 0.33
DHT + T	372 to 1349 ng/dL
SHBG	10.8 to 46.6 nmol/L
FSH	1.0 to 6.9 mIU/mL
LH	1.0 to 8.1 mIU/mL
E <sub>2</sub>	17.1 to 46.1 pg/mL

There is considerable variation in the half-life of testosterone reported in the literature, ranging from 10 to 100 minutes. Researchers do agree, however, that circulating testosterone has a diurnal variation in normal young men. Maximum levels occur at approximately 6:00 to 8:00 a.m. with levels declining throughout the day. Characteristic profiles have a maximum testosterone level of 720 ng/dL and a minimum level of 430 ng/dL. The physiological significance of this diurnal cycle, if any, however, is not clear.

### **B. Hypogonadal Men and Current Treatments for Hypogonadism**

Male hypogonadism results from a variety of patho-physiological conditions in which testosterone concentration is diminished below the normal range. The hypogonadic condition is sometimes linked with a number of physiological changes, such as diminished interest in sex, impotence, reduced lean body mass, decreased bone density, lowered mood, and energy levels.

Researchers generally classify hypogonadism into one of three types. Primary hypogonadism includes the testicular failure due to congenital or acquired anorchia, XYY Syndrome, XX males, Noonan's Syndrome, gonadal dysgenesis, Leydig cell tumors, undescended testes, varicocele, Sertoli-Cell-Only Syndrome, cryptorchidism, bilateral torsion, vanishing testis syndrome, orchiectomy, Klinefelter's Syndrome, chemotherapy, toxic damage from alcohol or heavy metals, and general disease (renal failure, liver cirrhosis, diabetes, myotonia dystrophica). Patients with primary hypogonadism show an intact feedback mechanism in that the low serum testosterone concentrations are associated with high FSH and LH concentrations. However, because of testicular or other failures, the high LH concentrations are not effective at stimulating testosterone production.

Secondary hypogonadism involves an idiopathic gonadotropin or LH-releasing hormone deficiency. This type of hypogonadism includes Kallman's Syndrome, Prader-Labhart-Willi's Syndrome, Laurence-Moon-Biedl's Syndrome, pituitary insufficiency/adenomas, Pasqualini's

10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100

Syndrome, hemochromatosis, hyperprolactinemia, or pituitary-hypothalamic injury from tumors, trauma, radiation, or obesity. Because patients with secondary hypogonadism do not demonstrate an intact feedback pathway, the lower testosterone concentrations are not associated with increased LH or FSH levels. Thus, these men have low testosterone serum levels but have gonadotropins in the normal to low range.

Third, hypogonadism may be age-related. Men experience a slow but continuous decline in average serum testosterone after approximately age 20 to 30 years. Researchers estimate that the decline is about 1-2% per year. Cross-sectional studies in men have found that the mean testosterone value at age 80 years is approximately 75% of that at age 30 years. Because the serum concentration of SHBG increases as men age, the fall in bioavailable and free testosterone is even greater than the fall in total testosterone. Researchers have estimated that approximately 50% of healthy men between the ages of 50 and 70 have levels of bioavailable testosterone that are below the lower normal limit. Moreover, as men age, the circadian rhythm of testosterone concentration is often muted, dampened, or completely lost. The major problem with aging appears to be within the hypothalamic-pituitary unit. For example, researchers have found that with aging, LH levels do not increase despite the low testosterone levels. Regardless of the cause, these untreated testosterone deficiencies in older men may lead to a variety of physiological changes, including sexual dysfunction, decreased libido, loss of muscle mass, decreased bone density, depressed mood, and decreased cognitive function. The net result is geriatric hypogonadism, or what is commonly referred to as “male menopause.”

Today, hypogonadism is the most common hormone deficiency in men, affecting 5 in every 1,000 men. At present, it is estimated that only five percent of the estimated four to five million American men of all ages with hypogonadism currently receive testosterone replacement

therapy. Thus, for years, researchers have investigated methods of delivering testosterone to men. These methods include intramuscular injections (43%), oral replacement (24%), pellet implants (23%), and transdermal patches (10%). A summary of these methods is shown in Table 2.

**Table 2: Mode of Application and Dosage of Various Testosterone Preparations**

Preparation	Route Of Application	Full Substitution Dose
<b>In Clinical Use</b>		
Testosterone enanthate	Intramuscular injection	200-250 mg every 2-3 weeks
Testosterone cypionate	Intramuscular injection	200 mg every 2 weeks
Testosterone undecanoate	Oral	2-4 capsules at 40 mg per day
Transdermal testosterone patch	Scrotal skin	1 membrane per day
Transdermal testosterone patch	Non-scrotal skin	1 or 2 systems per day
Testosterone implants	Implantation under the abdominal skin	3-6 implants of 200 mg every 6 months
<b>Under Development</b>		
Testosterone cyclodextrin	Sublingual	2.5-5.0 mg twice daily
Testosterone undecanoate	Intramuscular injection	1000 mg every 8-10 weeks
Testosterone buciclate	Intramuscular injection	1000 mg every 12-16 weeks
Testosterone microspheres	Intramuscular injection	315 mg for 11 weeks
<b>Obsolete</b>		
17 $\alpha$ -Methyltestosterone	Oral	25-50 mg per day
Fluoxymesterone	Sublingual	10-25 mg per day
	Oral	10-20 mg per day

As discussed below, all of the testosterone replacement methods currently employed suffer from one or more drawbacks, such as undesirable pharmacokinetic profiles or skin irritation. Thus, although the need for an effective testosterone replacement methodology has existed for decades, an alternative replacement therapy that overcomes these problems has never been developed. The present invention is directed to a 1% testosterone hydroalcoholic gel that overcomes the problems associated with current testosterone replacement methods.

### 1. Subdermal Pellet Implants

Subdermal implants have been used as a method of testosterone replacement since the 1940s. The implant is produced by melting crystalline testosterone into a cylindrical form.

Today, pellet implants are manufactured to contain either 100 mg (length 6 mm, surface area 117 mm<sup>2</sup>) or 200 mg of testosterone (length 12 mm, surface area 202 mm<sup>2</sup>). Patients receive dosages ranging from 100 to 1,200 mg, depending on the individual's requirements. The implants are inserted subcutaneously either by using a trocar and cannula or by open surgery into an area where there is relatively little movement. Frequently, the implant is placed in the lower abdominal wall or the buttock. Insertion is made under local anesthesia, and the wound is closed with an adhesive dressing or a fine suture.

Implants have several major drawbacks. First, implants require a surgical procedure which many hypogonadal men simply do not wish to endure. Second, implant therapy includes a risk of extrusion (8.5%), bleeding (2.3%), or infection (0.6%). Scarring is also a risk. Perhaps most important, the pharmacokinetic profile of testosterone pellet implant therapy fails to provide men with a suitable consistent testosterone level. In general, subdermal testosterone implants produce supra-physiologically high serum testosterone levels which slowly decline so that before the next injection subnormally low levels of testosterone are reached. For example, in one recent pharmacokinetic study, hypogonadal patients who received six implants (1,200 mg testosterone) showed an initial short-lived burst release of testosterone within the first two days after application. A stable plateau was then maintained over then next two months (day 2: 1,015 ng/dL; day 63: 990 ng/dL). Thereafter, the testosterone levels declined to baseline by day 300. DHT serum concentrations also rose significantly above the baseline, peaking at about 63 days after implementation and greatly exceeding the upper limit of the normal range. From day 21 to day 189, the DHT/T ratio was significantly increased. The pharmacokinetic profiles for testosterone, DHT, and DHT/T in this study are shown in FIG. 1. See Jockenhovel et al., *Pharmacokinetics and Pharmacodynamics of Subcutaneous Testosterone Implants in*



*Hypogonadal Men*, 45 CLINICAL ENDOCRINOLOGY 61-71 (1996). Other studies involving implants have reported similar undesirable pharmacokinetic profiles.

## 2. Injection of Testosterone Esters

Since the 1950s, researchers have experimented with the intermuscular depot injection of testosterone esters (such as enanthate, cypionate) to increase testosterone serum levels in hypogonadal men. More recent studies have involved injection of testosterone buciclate or testosterone undecanoate in an oil-based vehicle. Other researchers have injected testosterone microcapsule formulations.

Testosterone ester injection treatments suffer from many problems. Patients receiving injection therapy often complain that the delivery mechanism is painful and causes local skin reactions. In addition, testosterone microcapsule treatment requires two simultaneous intramuscular injections of a relatively large volume, which may be difficult to administer due to the high viscosity of the solution and the tendency to block the needle. Other men generally find testosterone injection therapy inconvenient because injection usually requires the patient to visit his physician every two to three weeks.

Equally important, injection-based testosterone replacement treatments still create an undesirable pharmacokinetic profile. The profile generally shows a supra-physiologic testosterone concentration during the first 24 to 48 hours followed by a gradual fall – often to sub-physiologic levels – over the next few weeks. These high serum testosterone levels, paralleled by increases in  $E_2$ , are also considered the reason for acne and gynecomastia occurring in some patients, and for polycythaemia, occasionally encountered especially in older patients using injectable testosterone esters. In the case of testosterone buciclate injections, the treatment barely provides normal androgen serum levels and the maximal increase of serum testosterone over baseline does not exceed 172 ng/dL (6 nmol/dL) on average. Because libido, potency,

mood, and energy are thought to fluctuate with the serum testosterone level, testosterone injections have largely been unsuccessful in influencing these variables. Thus, testosterone injection remains an undesirable testosterone replacement treatment method.

### 3. Oral/Sublingual/Buccal Preparations of Androgens

5 In the 1970s, researchers began using with oral, sublingual, or buccal preparations of androgens (such as fluoxymesterone, 17 $\alpha$ -methyl-testosterone or testosterone undecanoate) as a means for testosterone replacement. More recently, researchers have experimented with the sublingual administration of testosterone-hydroxypropyl-beta-cyclodextrin inclusion complexes. Predictably, both fluoxymesterone and methyl testosterone are 17-alkylated and thus associated with liver toxicity. Because these substances must first pass through the liver, they also produce an unfavorable effect on serum lipid profile, increasing LDL and decreasing HDL, and carbohydrate metabolism. While testosterone undecanoate has preferential absorption through the intestinal lymphatics, it has not been approved in the United States.

10 The pharmacokinetic profiles for oral, sublingual, and buccal delivery mechanisms are also undesirable because patients are subjected to super-physiologic testosterone levels followed by a quick return to the baseline. For example, one recent testing of a buccal preparation showed that patients obtained a peak serum hormone levels within 30 minutes after administration, with a mean serum testosterone concentration of 2,688 +/- 147 ng/dL and a return to baseline in 4 to 6 hours. See Dobs et al., *Pharmacokinetic Characteristics, Efficacy and Safety of Buccal*  
15 *Testosterone in Hypogonadal Males: A Pilot Study*, 83 J. CLINICAL ENDOCRINOLOGY & METABOLISM 33-39 (1998). To date, the ability these testosterone delivery mechanisms to alter physiological parameters (such muscle mass, muscle strength, bone resorption, urinary calcium excretion, or bone formation) is inconclusive. Likewise, researchers have postulated that super-

physiologic testosterone levels may not have any extra beneficial impact on mood parameters such as anger, nervousness, and irritability.

#### 4. Testosterone Transdermal Patches

The most recent testosterone delivery systems have involved transdermal patches. Currently, there are three patches used in the market: TESTODERM®, TESTODERM® TTS, and ANDRODERM®.

##### a. TESTODERM®

TESTODERM® (Alza Pharmaceuticals, Mountain View, CA) was the first testosterone-containing patch developed. The TESTODERM® patch is currently available in two sizes (40 or 60 cm<sup>2</sup>). The patch contains 10 or 15 mg of testosterone and delivers 4.0 mg or 6.0 mg of testosterone per day. TESTODERM® is placed on shaved scrotal skin, aided by application of heat for a few seconds from a hair dryer.

FIG. 2 shows a typical pharmacokinetic profile testosterone profile for both the 40 cm<sup>2</sup> and 60 cm<sup>2</sup> patch. Studies have also shown that after two to four weeks of continuous daily use, the average plasma concentration of DHT and DHT/T increased four to five times above normal. The high serum DHT levels are presumably caused by the increased metabolism of 5 $\alpha$ -reductase in the scrotal skin.

Several problems are associated with the TESTODERM® patch. Not surprisingly, many men simply do not like the unpleasant experience of dry-shaving the scrotal hair for optimal contact. In addition, patients may not be able to wear close-fitting underwear when undergoing treatment. Men frequently experience dislodgment of the patch, usually with exercise or hot weather. In many instances, men experience itching and/or swelling in the scrotal area. Finally, in a number of patients, there is an inability to achieve adequate serum hormone levels.

**b. TESTODERM® TTS**

The most recently developed non-scrotal patch is TESTODERM® TTS (Alza Pharmaceuticals, Mountain View, CA). It is an occlusive patch applied once daily to the arm, back, or upper buttocks. The system is comprised of a flexible backing of transparent polyester/ethylene-vinyl acetate copolymer film, a drug reservoir of testosterone, and an ethylene-vinyl acetate copolymer membrane coated with a layer of polyisobutylene adhesive formulation. A protective liner of silicone-coated polyester covers the adhesive surface.

Upon application, serum testosterone concentrations rise to a maximum at two to four hours and return toward baseline within two hours after system removal. Many men, however, are unable to obtain and/or sustain testosterone levels within the normal range. The pharmacokinetic parameters for testosterone concentrations are shown as follows:

**Table 3: TESTODERM® TTS Testosterone Parameters**

Parameters	Day 1	Day 5
C <sub>max</sub> (ng/dL)	482 ± 149	473 ± 148
T <sub>max</sub> (h)	3.9	3.0
C <sub>min</sub> (ng/dL)	164 ± 104	189 ± 86
T <sub>min</sub> (h)	0	0

The typical 24-hour steady state testosterone concentration achieved with TESTODERM® TTS patch is shown in FIG. 3.

Because of TESTODERM® patch is applied to the scrotal skin while the TESTODERM TTS® patch is applied to non-scrotal skin, the two patches provide different steady-state concentrations of the two major testosterone metabolites, DHT and E<sub>2</sub>:

**Table 4: Hormone Levels Using TESTODERM® and TESTODERM® TTS**

Hormone	Placebo	TESTODERM®	TESTODERM® TTS
DHT (ng/dL)	11	134	38
E <sub>2</sub> (pg/ml)	3.8	10	21.4

Likewise, in contrast to the scrotal patch, TESTODERM TTS<sup>®</sup> treatment creates a DHT/T ratio that is not different from that of a placebo treatment. Both systems, however, suffer from similar problems. In clinical studies, TESTODERM<sup>®</sup> TTS is associated with transient itching in 12% of patients, erythema in 3% of patients, and pruritus in 2% of patients. Moreover, in one 14-day study, 42% of patients reported three or more detachments, 33% of which occurred during exercise.

c. ANDRODERM<sup>®</sup>

ANDRODERM<sup>®</sup> (Watson Laboratories, Inc., Corona, CA) is a testosterone-containing patch applied to non-scrotal skin. The circular patch has a total surface area of 37 cm.<sup>2</sup> The patch consists of a liquid reservoir containing 12.2 mg of testosterone and a permeation-enhanced vehicle containing ethanol, water, monoglycerides, fatty acid esters, and gelling agents. The suggested dose of two patches, applied each night in a rotating manner on the back, abdomen, upper arm, or thigh, delivers 4.1 to 6.8 mg of testosterone.

The steady state pharmacokinetic profile of a clinical study involving ANDRODERM<sup>®</sup> is shown in FIG. 4. In general, upon repeated application of the ANDRODERM<sup>®</sup> patch, serum testosterone levels increase gradually for eight hours after each application and then remain at this plateau level for about another eight hours before declining.

In clinical trials, ANDRODERM<sup>®</sup> is associated with skin irritation in about a third of the patients, and 10% to 15% of subjects have been reported to discontinue the treatment because of chronic skin irritation. Preapplication of corticosteroid cream at the site of application of ANDRODERM<sup>®</sup> has been reported to decrease the incidence and severity of the skin irritation. A recent study, however, found that the incidence of skin reactions sufficiently noxious enough to interrupt therapy was as high as 52%. See Parker et al., *Experience with Transdermal*

(1999). The study reported:

Two-thirds of respondents found the Andropatch unsatisfactory. Patches were variously described as noisy, visually indiscrete, embarrassing, unpleasant to apply and remove, and generally to be socially unacceptable. They fell off in swimming pools and showers, attracted ribald comments from sporting partners, and left bald red marks over trunk and limbs. Dogs, wives, and children were distracted by noise of the patches with body movements. Those with poor mobility or manual dexterity (and several were over 70 years of age) found it difficult to remove packaging and apply patches dorsally.

#### d. Transdermal Patch Summary

In sum, the transdermal patch generally offers an improved pharmacokinetic profile compared to other currently used testosterone delivery mechanisms. However, as discussed above, the clinical and survey data shows that all of these patches suffer from significant drawbacks, such as buritis, burn-like blisters, and erythema. Moreover, one recent study has concluded that the adverse effects associated with transdermal patch systems are "substantially higher" than reported in clinical trials. See Parker, *supra*. Thus, the transdermal patch still remains an inadequate testosterone replacement therapy alternative for most men.

#### 5. DHT Gels

Researchers have recently begun investigating the application of DHT to the skin in a transdermal gel. However, the pharmacokinetics of a DHT-gel is markedly different from that of a testosterone gel. Application of DTH-gel results in decreased serum testosterone, E<sub>2</sub>, LH, and FSH levels. Thus, DHT gels are not effective at increasing testosterone levels in hypogonadal men.

Accordingly, there is a definite need for a testosterone formulation that safely and effectively provides an optimal and predictable pharmacokinetic profile.

## SUMMARY OF THE INVENTION

The foregoing problems are solved and a technical advance is achieved with the present invention. The present invention generally comprises a testosterone gel. Daily transdermal application of the gel in hypogonadal men results in a unique pharmacokinetic steady-state  
5 profile for testosterone. Long-term treatment further results in increased bone mineral density, enhanced libido, enhanced erectile frequency and satisfaction, increased positive mood, increased muscle strength, and improved body composition without significant skin irritation. The present invention is also directed to a unique method of administering the testosterone gel employing a packet having a polyethylene liner compatible with the components of the gel.

14111796V-2

## BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a graph of testosterone concentrations, DHT concentrations, and the DHT/T ratio for patients receiving a subdermal testosterone pellet implant over a period of 300 days after implantation.

5 FIG. 2 is a 24-hour testosterone pharmacokinetic profile for patients receiving the 40 cm<sup>2</sup> or 60 cm<sup>2</sup> TESTODERM<sup>®</sup> patch.

FIG. 3 is a 24-hour testosterone pharmacokinetic profile for patients receiving the TESTODERM<sup>®</sup> TTS patch.

10 FIG. 4 is a 24-hour testosterone pharmacokinetic profile for patients receiving the ANDRODERM<sup>®</sup> patch.

FIG. 5(a) is a graph showing the 24-hour testosterone pharmacokinetic profile for hypogonadal men prior to receiving 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

15 FIG. 5(b) is a graph showing the 24-hour testosterone pharmacokinetic profile for hypogonadal men on the first day of treatment with either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 5(c) is a graph showing the 24-hour testosterone pharmacokinetic profile for hypogonadal men on day 30 of treatment with either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel, or the testosterone patch (by initial treatment group).

20 FIG. 5(d) is a graph showing the 24-hour testosterone pharmacokinetic profile for hypogonadal men on day 90 of treatment with either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).



FIG. 5(e) is a graph showing the 24-hour testosterone pharmacokinetic profile for hypogonadal men on day 180 of treatment with either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by final treatment group).

5 FIG. 5(f) is a graph showing the 24-hour testosterone pharmacokinetic profile for hypogonadal men on day 0, 1, 30, 90, and 180 of treatment with 5.0 g/day of AndroGel<sup>®</sup>.

FIG. 5(g) is a graph showing the 24-hour testosterone pharmacokinetic profile for hypogonadal men on day 0, 1, 30, 90, and 180 of treatment with 10.0 g/day of AndroGel<sup>®</sup>.

FIG. 5(h) is a graph showing the 24-hour testosterone pharmacokinetic profile for hypogonadal men on day 0, 1, 30, 90, and 180 of treatment with the testosterone patch.

10 FIG. 6(a) is a graph showing the 24-hour free testosterone pharmacokinetic profile for hypogonadal men on day 1 of treatment with either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

15 FIG. 6(b) is a graph showing the 24-hour free testosterone pharmacokinetic profile for hypogonadal men on day 30 of treatment with either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 6(c) is a graph showing the 24-hour free testosterone pharmacokinetic profile for hypogonadal men on day 90 of treatment with either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

20 FIG. 6(d) is a graph showing the 24-hour free testosterone pharmacokinetic profile for hypogonadal men on day 180 of treatment with either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by final treatment group).

FIG. 6(e) is a graph showing the 24-hour free testosterone pharmacokinetic profile for hypogonadal men on day 0, 1, 30, 90, and 180 of treatment with 5.0 g/day of AndroGel<sup>®</sup>.

FIG. 6(f) is a graph showing the 24-hour free testosterone pharmacokinetic profile for hypogonadal men on day 0, 1, 30, 90, and 180 of treatment with 10.0 g/day of AndroGel®.

FIG. 6(g) is a graph showing the 24-hour free testosterone pharmacokinetic profile for hypogonadal men on day 0, 1, 30, 90, and 180 of treatment with the testosterone patch.

5 FIG. 7 is a graph showing the DHT concentrations on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel®, 10.0 g/day of AndroGel®, or the testosterone patch (by initial treatment group).

FIG. 8 is a graph showing the DHT/T ratio on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel®, 10.0 g/day of AndroGel®, or the testosterone patch (by initial treatment group).

FIG. 9 is a graph showing the total androgen concentrations (DHT +T) on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel®, 10.0 g/day of AndroGel®, or the testosterone patch (by initial treatment group).

FIG. 10 is a graph showing the E<sub>2</sub> concentrations on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel®, 10.0 g/day of AndroGel®, or the testosterone patch (by initial treatment group).

FIG. 11 is a graph showing the SHBG concentrations on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel®, 10.0 g/day of AndroGel®, or the testosterone patch (by initial treatment group).

20 FIG. 12(a) is a graph showing the FSH concentrations on days 0 through 180 for men having primary hypogonadism and receiving either 5.0 g/day of AndroGel®, 10.0 g/day of AndroGel®, or the testosterone patch (by initial treatment group).

FIG. 12(b) is a graph showing the FSH concentrations on days 0 through 180 for men having secondary hypogonadism and receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 12(c) is a graph showing the FSH concentrations on days 0 through 180 for men having age-associated hypogonadism and receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 12(d) is a graph showing the FSH concentrations on days 0 through 180 for men having hypogonadism of an unknown origin and receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 13(a) is a graph showing the LH concentrations on days 0 through 180 for men having primary hypogonadism and receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 13(b) is a graph showing the LH concentrations on days 0 through 180 for men having secondary hypogonadism and receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 13(c) is a graph showing the LH concentrations on days 0 through 180 for men having age-associated hypogonadism and receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 13(d) is a graph showing the LH concentrations on days 0 through 180 for men having hypogonadism of an unknown origin and receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 14(a) is a bar graph showing the change in hip BMD for hypogonadal men after 180 days of treatment with 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 14(b) is a bar graph showing the change in spine BMD for hypogonadal men after 180 days of treatment with 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch

FIG. 15 is a graph showing PTH concentrations on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 16 is a graph showing SALP concentrations on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 17 is a graph showing the osteocalcin concentrations on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 18 is a graph showing the type I procollagen concentrations on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 19 is a graph showing the N-telopeptide/Cr ratio on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 20 is a graph showing the Ca/Cr ratio on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch (by initial treatment group).

FIG. 21(a) is a graph showing sexual motivation scores on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 21(b) is a graph showing overall sexual desire scores on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 21(c) is a graph showing sexual enjoyment (with a partner) scores on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 22(a) is a graph showing sexual performance scores on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 22(b) is a graph showing erection satisfaction performance scores on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 22(c) is a graph showing percent erection scores on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 23(a) is a graph showing positive mood scores on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

5 FIG. 23(b) is a graph showing negative mood scores on days 0 through 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 24(a) is a bar graph showing the change in leg strength on days 90 and 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

10 FIG. 24(b) is a bar graph showing the change in arm strength on days 90 and 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

15 FIG. 25(a) is a bar graph showing the change in total body mass on days 90 and 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 25(b) is a bar graph showing the change in lean body mass on days 90 and 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

20 FIG. 25(c) is a bar graph showing the change in fat mass on days 90 and 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 25(d) is a bar graph showing the change in percent body fat on days 90 and 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

FIG. 25(d) is a bar graph showing the change in percent body fat on days 90 and 180 for hypogonadal men receiving either 5.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day 10.0 g/day of AndroGel<sup>®</sup>, or the testosterone patch.

## DETAILED DESCRIPTION OF THE INVENTION

While the present invention may be embodied in many different forms, several specific embodiments are discussed herein with the understanding that the present disclosure is to be considered only as an exemplification of the principles of the invention, and it is not intended to  
5 limit the invention to the embodiments illustrated.

The present invention is directed to a pharmaceutical composition for percutaneous administration comprising at least one active pharmaceutical ingredient (e.g., testosterone) in a hydroalcoholic gel. In a broad aspect of the invention, the active ingredients employed in the composition may include anabolic steroids such as androisoxazole, bolasterone, clostebol, ethylestrenol, formyldienolone, 4-hydroxy-19-nortestosterone, methenolone, methyltrienolone, nandrolone, oxymesterone, quinbolone, stenbolone, trenbolone; androgenic steroids such as boldenone, fluoxymesterone, mestanolone, mesterolone, methandrostenolone, 17-methyltestosterone, 17  $\alpha$ -methyl-testosterone 3-cyclopentyl enol ether, norethandrolone, normethandrone, oxandrolone, oxymetholone, prasterone, stanlolone, stanozolol, dihydrotestosterone, testosterone; and progestogens such as anagestone, chlormadinone acetate, delmadinone acetate, demegestone, dimethisterone, dihydrogesterone, ethinylestrenol, ethisterone, ethynodiol, ethynodiol diacetate, flurogestone acetate, gestodene, gestonorone caproate, haloprogestosterone, 17-hydroxy-16-methylene-progesterone, 17  $\beta$ -hydroxyprogesterone, 17  $\alpha$ -hydroxyprogesterone caproate, medrogestone, medroxyprogesterone, megestrol acetate,  
10 melengestrol, norethindrone, norethindrone acetate, norethynodrel, norgesterone, norgestimate, norgestrel, norgestrienone, 19-norprogesterone, norvinisterone, pentagestrone, progesterone, promegestone, quingestrone, and trengestone; and all enantiomers, isomers and derivatives of  
20



these compounds. (Based upon the list provided in The Merck Index, Merck & Co. Rahway, N.J. (1998)).

In addition to the active ingredient, the gel comprises one or more lower alcohols, such as ethanol or isopropanol; a penetration enhancing agent; a thickener; and water. Additionally, the present invention may optionally include salts, emollients, stabilizers, antimicrobials, fragrances, and propellants.

A "penetration enhancer" is an agent known to accelerate the delivery of the drug through the skin. These agents also have been referred to as accelerants, adjuvants, and sorption promoters, and are collectively referred to herein as "enhancers." This class of agents includes those with diverse mechanisms of action including those which have the function of improving the solubility and diffusibility of the drug, and those which improve percutaneous absorption by changing the ability of the stratum corneum to retain moisture, softening the skin, improving the skin's permeability, acting as penetration assistants or hair-follicle openers or changing the state of the skin such as the boundary layer.

The penetration enhancer of the present invention is a functional derivative of a fatty acid, which includes isosteric modifications of fatty acids or non-acidic derivatives of the carboxylic functional group of a fatty acid or isosteric modifications thereof. In one embodiment, the functional derivative of a fatty acid is an unsaturated alkanolic acid in which the —COOH group is substituted with a functional derivative thereof, such as alcohols, polyols, amides and substituted derivatives thereof. The term "fatty acid" means a fatty acid that has four (4) to twenty-four (24) carbon atoms. Non-limiting examples of penetration enhancers include C8-C22 fatty acids such as isostearic acid, octanoic acid, and oleic acid; C8-C22 fatty alcohols such as oleyl alcohol and lauryl alcohol; lower alkyl esters of C8-C22 fatty acids such as ethyl

oleate, isopropyl myristate, butyl stearate, and methyl laurate; di(lower)alkyl esters of C6-C8 diacids such as diisopropyl adipate; monoglycerides of C8-C22 fatty acids such as glyceryl monolaurate; tetrahydrofurfuryl alcohol polyethylene glycol ether; polyethylene glycol, propylene glycol; 2-(2-ethoxyethoxy)ethanol; diethylene glycol monomethyl ether; alkylaryl ethers of polyethylene oxide; polyethylene oxide monomethyl ethers; polyethylene oxide dimethyl ethers; dimethyl sulfoxide; glycerol; ethyl acetate; acetoacetic ester; N-alkylpyrrolidone; and terpenes.

The thickeners used herein may include anionic polymers such as polyacrylic acid (CARBOPOL® by B.F. Goodrich Specialty Polymers and Chemicals Division of Cleveland, Ohio), carboxymethylcellulose and the like. Additional thickeners, enhancers and adjuvants may generally be found in United States Pharmacopeia/National Formulary (2000); Remington's The Science and Practice of Pharmacy, Meade Publishing Co.

The amount of drug to be incorporated in the composition varies depending on the particular drug, the desired therapeutic effect, and the time span for which the gel is to provide a therapeutic effect. The composition is used in a "pharmacologically effective amount." This means that the concentration of the drug is such that in the composition it results in a therapeutic level of drug delivered over the term that the gel is to be used. Such delivery is dependent on a number of variables including the drug, the form of drug, the time period for which the individual dosage unit is to be used, the flux rate of the drug from the gel, surface area of application site, etc. The amount of drug necessary can be experimentally determined based on the flux rate of the drug through the gel, and through the skin when used with and without enhancers.

One such testosterone gel has only recently been made available in the United States under the trademark AndroGel® by Unimed Pharmaceuticals, Inc., Deerfield, Illinois, one of the assignees of this application. In one embodiment, the gel is comprised of the following substances in approximate amounts:

**Table 5: Composition of AndroGel®**

SUBSTANCE	AMOUNT (w/w) PER 100g OF GEL
Testosterone	1.0 g
Carbopol 980	0.90 g
Isopropyl myristate	0.50 g
0.1 N NaOH	4.72 g
Ethanol (95% w/w)	72.5 g*
Purified water (qsf)	100 g

\*Corresponding to 67 g of ethanol.

One skilled in the art will appreciate that the constituents of this formulation may be varied in amounts yet continue to be within the spirit and scope of the present invention. For example, the composition may contain about 0.1 to about 10.0 g of testosterone, about 0.1 to about 5.0 g Carbopol, about 0.1 to about 5.0 g isopropyl myristate, and about 30.0 to about 98.0 g ethanol.

A therapeutically effective amount of the gel is rubbed onto a given area of skin by the user. The combination of the lipophilic testosterone with the hydroalcoholic gel helps drive the testosterone in to the outer layers of the skin where it is absorbed and then slowly released into the blood stream. As demonstrated by the data presented herein, the administration of the gel of the present invention has a sustained effect.

Toxicity and therapeutic efficacy of the active ingredients can be determined by standard pharmaceutical procedures, *e.g.*, for determining LD<sub>50</sub> (the dose lethal to 50% of the population) and the ED<sub>50</sub> (the dose therapeutically effective in 50% of the population). The dose ratio

between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio  $LD_{50}/ED_{50}$ . Compounds which exhibit large therapeutic induces are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The term "treatment" as used herein refers to any treatment of a human condition or disease and includes: (1) preventing the disease or condition from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it, (2) inhibiting the disease or condition, *i.e.*, arresting its development, (3) relieving the disease or condition, *i.e.*, causing regression of the condition, or (4) relieving the conditions caused by the disease, *i.e.*, stopping the symptoms of the disease.

Although the examples of the present invention involve the treatment of disorders associated with hypogonadal men, the composition and method of the present invention may be used to treat these disorders in humans and animals of any kind, such as dogs, pigs, sheep, horses, cows, cats, zoo animals, and other commercially bred farm animals.

The present invention is further illustrated by the following examples, which should not be construed as limiting in any way. The contents of all cited references throughout this application are hereby expressly incorporated by reference. The practice of the present invention will employ, unless otherwise indicated, conventional techniques of pharmacology and pharmaceuticals, which are within the skill of the art.

## EXAMPLES

### **Example 1: Treatment of Hypogonadism in Male Subjects**

One embodiment of the present invention involves the transdermal application of AndroGel® as a method of treating male hypogonadism. As demonstrated below, application of the gel results in a unique pharmacokinetic profile for testosterone, as well as concomitant modulation of several other sex hormones. Application of the testosterone gel to hypogonadal male subjects also results in: (1) increased bone mineral density, (2) enhanced libido, (3) enhanced erectile capability and satisfaction, (4) increased positive mood, (5) increased muscle strength, and (6) better body composition, such increased total body lean mass and decreased total body fat mass. Moreover, the gel is not associated with significant skin irritation.

#### ***Methods***

In this example, hypogonadal men were recruited and studied in 16 centers in the United States. The patients were between 19 and 68 years and had single morning serum testosterone levels at screening of less than or equal to 300 ng/dL (10.4 nmol/L ). A total of 227 patients were enrolled: 73, 78, and 76 were randomized to receive 5.0 g/day of AndroGel® (delivering 50 mg/day of testosterone to the skin of which about 10% or 5 mg is absorbed), 10.0 g/day of AndroGel® (delivering 100 mg/day of testosterone to the skin of which about 10% or 10 mg is absorbed), or the ANDRODERM® testosterone patch ("T patch") (delivering 50 mg/day of testosterone), respectively.

As shown in the following table, there were no significant group-associated differences of the patients' characteristics at baseline.

**Table 6. Baseline Characteristics of the Hypogonadal Men**

<b>Treatment Group</b>	<b>T patch</b>	<b>AndroGel® (5.0 g/day)</b>	<b>AndroGel® (10.0 g/day)</b>
No of subjects enrolled	76	73	78

Treatment Group	T patch	AndroGel® (5.0 g/day)	AndroGel® (10.0 g/day)
Age (years)	51.1	51.3	51.0
Range (years)	28-67	23-67	19-68
Height (cm)	179.3 ± 0.9	175.8 ± 0.8	178.6 ± 0.8
Weight (kg)	92.7 ± 1.6	90.5 ± 1.8	91.6 ± 1.5
Serum testosterone (nmol/L)	6.40 ± 0.41	6.44 ± 0.39	6.49 ± 0.37
Causes of hypogonadism			
Primary hypogonadism	34	26	34
Klinefelter's Syndrome	9	5	8
Post Orchidectomy/Anorchia	2	1	3
Primary Testicular Failure	23	20	23
Secondary hypogonadism	15	17	12
Kallman's Syndrome	2	2	0
Hypothalimic Pituitary Disorder	6	6	3
Pituitary Tumor	7	9	9
Aging	6	13	6
Not classified	21	17	26
Years diagnosed	5.8 ± 1.1	4.4 ± 0.9	5.7 ± 1.24
Number previously treated with testosterone	50 (65.8%)	38 (52.1%)	46 (59.0%)
Type of Previous Hormonal Treatment			
Intramuscular injections	26	20	28
Transdermal patch	12	7	8
All others	12	11	10
Duration of treatment (years)	5.8 ± 1.0	5.4 ± 0.8	4.6 ± 80.7

Forty-one percent (93/227) of the subjects had not received prior testosterone replacement therapy. Previously treated hypogonadal men were withdrawn from testosterone ester injection for at least six weeks and oral or transdermal androgens for four weeks before the screening visit. Aside from the hypogonadism, the subjects were in good health as evidenced by medical history, physical examination, complete blood count, urinalysis, and serum biochemistry. If the subjects were on lipid-lowering agents or tranquilizers, the doses were stabilized for at least three months prior to enrollment. Less than 5% of the subjects were taking supplemental calcium or vitamin D during the study. The subjects had no history of chronic medical illness, alcohol or drug abuse. They had a normal rectal examination, a PSA level of less than 4 ng/mL, and a urine flow rate of 12 mL/s or greater. Patients were excluded if they

had a generalized skin disease that might affect the testosterone absorption or prior history of skin irritability with ANDRODERM<sup>®</sup> patch. Subjects weighing less than 80% or over 140% of their ideal body weight were also excluded.

The randomized, multi-center, parallel study compared two doses of AndroGel<sup>®</sup> with the ANDRODERM<sup>®</sup> testosterone patch. The study was double-blind with respect to the AndroGel<sup>®</sup> dose and open-labeled for the testosterone patch group. For the first three months of the study (days 1 to 90), the subjects were randomized to receive 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, or two non-scrotal patches. In the following three months (days 91 to 180), the subjects were administered one of the following treatments: 5.0 g/day of AndroGel<sup>®</sup>, 10.0 g/day of AndroGel<sup>®</sup>, 7.5 g/day of AndroGel<sup>®</sup>, or two non-scrotal patches. Patients who were applying AndroGel<sup>®</sup> had a single, pre-application serum testosterone measured on day 60 and, if the levels were within the normal range of 300 to 1,000 ng/dL (10.4 to 34.7 nmol/L ), then they remained on their original dose. Patients with testosterone levels less than 300 ng/dL and who were originally assigned to apply 5.0 g/day of AndroGel<sup>®</sup> and those with testosterone levels more than 1,000 ng/dL who had received 10.0 g/day of AndroGel<sup>®</sup> were then reassigned to administer 7.5 g/day of AndroGel<sup>®</sup> for days 91 to 180.

Accordingly, at 90 days, dose adjustments were made in the AndroGel<sup>®</sup> groups based on the pre-application serum testosterone levels on day 60. Twenty subjects in the 5.0 g/day AndroGel<sup>®</sup> group had the dose increased to 7.5 g/day. Twenty patients in the 10.0 g/day AndroGel<sup>®</sup> group had the AndroGel<sup>®</sup> dose reduced to 7.5 g/day. There were three patients in the testosterone patch group who were switched to 5.0 g/day AndroGel<sup>®</sup> because of patch intolerance. One 10.0 g/day AndroGel<sup>®</sup> subject was adjusted to receive 5.0 g/day and one 5.0 g/day AndroGel<sup>®</sup> subject had the dose adjusted to 2.5 g/day. The number of subjects

enrolled into day 91 to 180 of the study thus consisted of 51 receiving 5.0 g/day of AndroGel<sup>®</sup>, 40 receiving 7.5 g/day of AndroGel<sup>®</sup>, 52 receiving 10.0 g/day of AndroGel<sup>®</sup>, and 52 continuing on the ANDRODERM<sup>®</sup> patch. The treatment groups in this example may thus be characterized in two ways, either by “initial” or by the “final” treatment group.

5        Subjects returned to the study center on days 0, 30, 60, 90, 120, 150, and 180 for a clinical examination, skin irritation and adverse event assessments. Fasting blood samples for calcium, inorganic phosphorus, parathyroid hormone (“PTH”), osteocalcin, type I procollagen, and skeletal specific alkaline phosphatase (“SALP”) were collected on days 0, 30, 90, 120, and 180. In addition, a fasting two-hour timed urine collection for urine creatinine, calcium, and type 1 collagen cross-linked N-telopeptides (“N-telopeptide”) were collected on days 0, 30, 90, 120, and 180. Other tests performed were as follows:

(1) *Hematology*: hemoglobin, hematocrit, red blood cell count, platelets, white blood cell counts with differential analysis (neutrophils, lymphocytes, monocytes, eosinophils, and basophils);

(2) *Chemistry*: alkaline phosphatase, alanine aminotransferase, serum glutamic pyruvic transaminase (“ALT/SGPT”), aspartate aminotransferase/serum glutamin axaloacetic transaminase (“AST/SGOT”), total bilirubin, creatinine, glucose, and electrolytes (sodium, potassium, chloride, bicarbonate, calcium, and inorganic phosphorus);

(3) *Lipids*: total cholesterol, high-density lipoprotein (“HDL”), low-density lipoprotein (“LDL”), and triglycerides;

(4) *Urinalysis*: color, appearance, specific gravity, pH, protein, glucose, ketones, blood, bilirubin, and nitrites; and

(5) *Other*: PSA (screening days 90-180), prolactin (screening), and testosterone (screening) including electrolytes, glucose, renal, and liver function tests and lipid profile, were performed at all clinic visits. Bone mineral density (“BMD”) was analyzed at day 0 and day 180.



**A. AndroGel® and ANDRODERM® patch**

Approximately 250 g of AndroGel® was packaged in multidose glass bottles that delivered 2.25 g of the gel for each actuation of the pump. Patients assigned to apply 5.0 g/day of AndroGel® testosterone were given one bottle of AndroGel® and one bottle of placebo gel (containing vehicle but no testosterone), while those assigned to receive 10.0 g/day of AndroGel® were dispensed two bottles of the active AndroGel®. The patients were then instructed to apply the bottle contents to the right and left upper arms/shoulders and to the right and left sides of the abdomen on an alternate basis. For example, on the first day of the study, patients applied two actuations from one bottle, one each to the left and right upper arm/shoulder, and two actuations from the second bottle, one each to the left and right abdomen. On the following day of treatment, the applications were reversed. Alternate application sites continued throughout the study. After application of the gel to the skin, the gel dried within a few minutes. Patients washed their hands thoroughly with soap and water immediately after gel application.

The 7.5 g/day AndroGel® group received their dose in an open-label fashion. After 90 days, for the subjects titrated to the AndroGel® 7.5 g/day dose, the patients were supplied with three bottles, one containing placebo and the other two AndroGel®. The subjects were instructed to apply one actuation from the placebo bottle and three actuations from a AndroGel® bottle to four different sites of the body as above. The sites were rotated each day taking the same sequence as described above.

ANDRODERM® testosterone patches each delivering 2.5 mg/day of testosterone were provided to about one-third of the patients in the study. These patients were instructed to apply two testosterone patches to a clean, dry area of skin on the back, abdomen, upper arms, or thighs

once per day. Application sites were rotated with approximately seven days interval between applications to the same site.

On study days when the patients were evaluated, the gel/patches were applied following pre-dose evaluations. On the remaining days, the testosterone gel or patches were applied at approximately 8:00 a.m. for 180 days.

## B. Study Method and Results

### 1. Hormone Pharmacokinetics

On days 0, 1, 30, 90, and 180, the patients had multiple blood samples for testosterone and free testosterone measurements at 30, 15 and 0 minutes before and 2, 4, 8, 12, 16, and 24 hours after AndroGel® or patch application. In addition, subjects returned on days 60, 120, and 150 for a single blood sampling prior to application of the gel or patch. Serum DHT, E<sub>2</sub>, FSH, LH and SHBG were measured on samples collected before gel application on days 0, 30, 60, 90, 120, 150, and 180. Sera for all hormones were stored frozen at -20 °C until assay. All samples for a patient for each hormone were measured in the same assay whenever possible. The hormone assays were then measured at the Endocrine Research Laboratory of the UCLA-Harbor Medical Center.

The following table summarizes the pharmacokinetic parameters were measured for each patient:

**Table 7: Pharmacokinetic Parameters**

AUC <sub>0-24</sub>	area under the curve from 0 to 24 hours, determined using the linear trapezoidal rule.
C <sub>base</sub> or C <sub>0</sub>	Baseline concentration
C <sub>avg</sub>	time-averaged concentration over the 24-hour dosing interval determined by AUC <sub>0-24</sub> /24
C <sub>max</sub>	maximum concentration during the 24-hour dosing interval

$C_{\min}$	minimum concentration during the 24-hour dosing interval
$T_{\max}$	time at which $C_{\max}$ occurred
$T_{\min}$	time at which $C_{\min}$ occurred
Fluctuation Index	extent of variation in the serum concentration over the course of a single day, calculated as $(C_{\max} - C_{\min})/C_{\text{avg}}$
Accumulation ratio	increase in the daily drug exposure with continued dosing, calculated as the ratio of the AUC at steady on a particular day over the AUC on day 1 (e.g., $\text{AUC}_{\text{day 30}}/\text{AUC}_{\text{day 1}}$ )
Net $\text{AUC}_{0-24}$	$\text{AUC}_{0-24}$ on days 30, 90, 180 - $\text{AUC}_{0-24}$ on day 0

#### a. Testosterone Pharmacokinetics

##### (1) Methods

Serum testosterone levels were measured after extraction with ethylacetate and hexane by a specific radioimmunoassay ("RIA") using reagents from ICN (Costa Mesa, CA). The cross reactivities of the antiserum used in the testosterone RIA were 2.0% for DHT, 2.3% for androstenedione, 0.8% for 3- $\beta$ -androstenediol, 0.6% for etiocholanolone and less than 0.01% for all other steroids tested. The lower limit of quantitation ("LLQ") for serum testosterone measured by this assay was 25 ng/dL (0.87 nmol/L). The mean accuracy of the testosterone assay, determined by spiking steroid free serum with varying amounts of testosterone (0.9 nmol/L to 52 nmol/L), was 104% and ranged from 92% to 117%. The intra-assay and inter-assay coefficients of the testosterone assay were 7.3 and 11.1%, respectively, at the normal adult male range. In normal adult men, testosterone concentrations range from 298 to 1,043 ng/dL (10.33 to 36.17 nmol/L) as determined at the UCLA-Harbor Medical Center.

##### (2) Baseline Concentration

As shown in Table 8 and FIG. 5(a), at baseline, the average serum testosterone concentrations over 24 hours ( $C_{\text{avg}}$ ) were similar in the groups and below the adult normal range. Moreover the variations of the serum concentration (based on maximum and minimum

concentrations during the 24-hour period,  $C_{max}$  and  $C_{min}$ , respectively) during the day were also similar in the three groups. FIG. 5(a) shows that the mean testosterone levels had a the maximum level between 8 to 10 a.m. (i.e., at 0 to 2 hours) and the minimum 8 to 12 hours later, demonstrating a mild diurnal variation of serum testosterone. About one-third of the patients in each group had  $C_{avg}$  within the lower normal adult male range on day 0 (24/73 for the 5.0 g/day AndroGel<sup>®</sup> group, 26/78 for the 10.0 g/day AndroGel<sup>®</sup> group, and 25/76 for testosterone patch group). All except three of the subjects met the enrollment criterion of serum testosterone less than 300 ng/dL (10.4 nmol/L) on admission.

**Table 8(a): Baseline Pharmacokinetic Parameters  
by Initial Treatment Group (Mean  $\pm$  SD)**

	5.0 g/day T-Gel	10.0 g/day T-gel	T-patch
N	73	78	76
$C_{avg}$ (ng/dL)	237 $\pm$ 130	248 $\pm$ 140	237 $\pm$ 139
$C_{max}$ (ng/dL)	328 $\pm$ 178	333 $\pm$ 194	314 $\pm$ 179
$T_{max}$ * (hr)	4.0 (0.0-24.5)	7.9 (0.0-24.7)	4.0 (0.0-24.3)
$C_{min}$ (ng/dL)	175 $\pm$ 104	188 $\pm$ 112	181 $\pm$ 112
$T_{min}$ * (hr)	8.01 (0.0-24.1)	8.0 (0.0-24.0)	8.0 (0.0-23.9)
Fluc Index (ratio)	0.627 $\pm$ 0.479	0.556 $\pm$ 0.384	0.576 $\pm$ 0.341

\*Median (Range\*)

**Table 8(b): Baseline Testosterone Pharmacokinetic Parameters  
by Final Treatment Group (Mean  $\pm$  SD)**

	Doses Received During Initial => Extended Treatment Phases				
	5.0 g/day T-gel	5.0 => 7.5 g/day T-gel	10.0 => 7.5 g/day T-gel	10.0 g/day T-gel	T-patch
N	53	20	20	58	76
$C_{avg}$ (ng/dL)	247 $\pm$ 137	212 $\pm$ 109	282 $\pm$ 157	236 $\pm$ 133	237 $\pm$ 140
$C_{max}$ (ng/dL)	333 $\pm$ 180	313 $\pm$ 174	408 $\pm$ 241	307 $\pm$ 170	314 $\pm$ 179
$T_{max}$ * (hr)	4.0 (0.0-24.5)	4.0 (0.0-24.0)	19.7 (0.0-24.3)	4.0 (0.0-24.7)	4.0 (0.0-24.3)
$C_{min}$ (ng/dL)	185 $\pm$ 111	150 $\pm$ 80	206 $\pm$ 130	182 $\pm$ 106	181 $\pm$ 112
$T_{min}$ * (hr)	8.0 (0.0-24.1)	11.9 (0.0-24.0)	8.0 (0.0-23.3)	8.0 (0.0-24.0)	8.0 (0.0-23.9)
Fluc Index (ratio)	0.600 $\pm$ 0.471	0.699 $\pm$ 0.503	0.678 $\pm$ 0.580	0.514 $\pm$ 0.284	0.576 $\pm$ 0.341

\*Median (range)

### (3) Day 1

FIG. 5(b) and Tables 8(c)-(d) show the pharmacokinetic profile for all three initial treatment groups after the first application of transdermal testosterone. In general, treatment

with AndroGel® and the testosterone patch produced increases in testosterone concentrations sufficiently large to bring the patients into the normal range in just a few hours. However, even on day 1, the pharmacokinetic profiles were markedly different in the AndroGel® and patch groups. Serum testosterone rose most rapidly in the testosterone patch group reaching a maximum concentration ( $C_{max}$ ) at about 12 hours ( $T_{max}$ ). In contrast, serum testosterone rose steadily to the normal range after AndroGel® application with  $C_{max}$  levels achieved by 22 and 16 hours in the 5.0 g/day AndroGel® group and the 10.0 g/day AndroGel® group, respectively.

**Table 8(c): Testosterone Pharmacokinetic Parameters on Day 1 by Initial Treatment Group (Mean  $\pm$  SD)**

	5.0 g/day T-Gel	10.0 g/day T-gel	T-patch
N	73	76	74
$C_{avg}$ (ng/dL)	398 $\pm$ 156	514 $\pm$ 227	482 $\pm$ 204
$C_{max}$ (ng/dL)	560 $\pm$ 269	748 $\pm$ 349	645 $\pm$ 280
$T_{max}^*$ (hr)	22.1 (0.0-25.3)	16.0 (0.0-24.3)	11.8 (1.8-24.0)
$C_{min}$ (ng/dL)	228 $\pm$ 122	250 $\pm$ 143	232 $\pm$ 132
$T_{min}^*$ (hr)	1.9 (0.0-24.0)	0.0 (0.0-24.2)	1.5 (0.0-24.0)

\*Median (Range)

**Table 8(d): Testosterone Pharmacokinetic Parameters on Day 1 by Final Treatment Group (Mean  $\pm$  SD)**

	Doses Received During Initial $\Rightarrow$ Extended Treatment Phases				
	5.0 g/day T-gel	5.0 $\Rightarrow$ 7.5 g/day T-gel	10.0 $\Rightarrow$ 7.5 g/day T-gel	10.0 g/day T-gel	T-patch
N	53	20	19	57	74
$C_{avg}$ (ng/dL)	411 $\pm$ 160	363 $\pm$ 143	554 $\pm$ 243	500 $\pm$ 223	482 $\pm$ 204
$C_{max}$ (ng/dL)	573 $\pm$ 285	525 $\pm$ 223	819 $\pm$ 359	724 $\pm$ 346	645 $\pm$ 280
$T_{max}^*$ (hr)	22.1 (0.0-25.3)	19.5 (1.8-24.3)	15.7 (3.9-24.0)	23.0 (0.0-24.3)	11.8 (1.8-24.0)
$C_{min}$ (ng/dL)	237 $\pm$ 125	204 $\pm$ 112	265 $\pm$ 154	245 $\pm$ 140	232 $\pm$ 132
$T_{min}^*$ (hr)	1.8 (0.0-24.0)	3.5 (0.0-24.0)	1.9 (0.0-24.2)	0.0 (0.0-23.8)	1.5 (0.0-24.0)
Fluc Index (ratio)	0.600 $\pm$ 0.471	0.699 $\pm$ 0.503	0.678 $\pm$ 0.580	0.514 $\pm$ 0.284	0.576 $\pm$ 0.341

\*Median (range)

#### (4) Days 30, 90, and 180

FIGS. 5(c) and 5(d) show the unique 24-hour pharmacokinetic profile of AndroGel®-treated patients on days 30 and 90. In the AndroGel® groups, serum testosterone levels showed small and variable increases shortly after dosing. The levels then returned to a relatively constant level. In contrast, in the testosterone patch group, patients exhibited a rise over the first

8 to 12 hours, a plateau for another 8 hours, and then a decline to the baseline of the prior day. Further, after gel application on both days 30 and 90, the  $C_{avg}$  in the 10.0 g/day AndroGel<sup>®</sup> group was 1.4 fold higher than in the 5.0 g/day AndroGel<sup>®</sup> group and 1.9 fold higher than the testosterone patch group. The testosterone patch group also had a  $C_{min}$  substantially below the lower limit of the normal range. On day 30, the accumulation ratio was 0.94 for testosterone patch group, showing no accumulation. The accumulation ratios at 1.54 and 1.9 were significantly higher in the 5.0 g/day AndroGel<sup>®</sup> group and 10.0 g/day AndroGel<sup>®</sup> group, respectively. The differences in accumulation ratio among the groups persisted on day 90. This data indicates that the AndroGel<sup>®</sup> preparations had a longer effective half-life than testosterone patch.

FIG. 5(e) shows the 24-hour pharmacokinetic profile for the treatment groups on day 180. In general, as Table 8(e) shows, the serum testosterone concentrations achieved and the pharmacokinetic parameters were similar to those on days 30 and 90 in those patients who continued on their initial randomized treatment groups. Table 8(f) shows that the patients titrated to the 7.5 g/day AndroGel<sup>®</sup> group were not homogeneous. The patients that were previously in the 10.0 g/day group tended to have higher serum testosterone levels than those previously receiving 5.0 g/day. On day 180, the  $C_{avg}$  in the patients in the 10.0 g/day group who converted to 7.5 g/day on day 90 was 744 ng/dL, which was 1.7 fold higher than the  $C_{avg}$  of 450 ng/dL in the patients titrated to 7.5 g/day from 5.0 g/day. Despite adjusting the dose up by 2.5 g/day in the 5.0 to 7.5 g/day group, the  $C_{avg}$  remained lower than those remaining in the 5.0 g/day group. In the 10.0 to 7.5 g/day group, the  $C_{avg}$  became similar to those achieved by patients remaining in the 10.0 g/day group without dose titration. These results suggest that many of the under-responders may actually be poorly compliant patients. For example, if a

patient does not apply AndroGel® properly (e.g., preferentially from the placebo container or shortly before bathing), then increasing the dose will not provide any added benefit.

FIGS. 5(f)-(h) compare the pharmacokinetic profiles for the 5.0 g/day AndroGel® group, the 10.0 AndroGel® g/day group, and the testosterone patch group at days 0, 1, 30, 90, and 180, respectively. In general, the mean serum testosterone levels in the testosterone patch group remained at the lower limit of the normal range throughout the treatment period. In contrast, the mean serum testosterone levels remained at about 490-570 ng/dL for the 5.0 g/day AndroGel® group and about 630-860 ng/dL AndroGel® for the 10.0 g/day group.

**Table 8(e): Testosterone Pharmacokinetic Parameters on Day 1  
by Initial Treatment Group (Mean ± SD)**

	5.0 g/day T-Gel	10.0 g/day T-gel	T-patch
<b>Day 30</b>	N = 66	N = 74	N = 70
C <sub>avg</sub> (ng/dL)	566 ± 262	792 ± 294	419 ± 163
C <sub>max</sub> (ng/dL)	876 ± 466	1200 ± 482	576 ± 223
T <sub>max</sub> * (hr)	7.9 (0.0-24.0)	7.8 (0.0-24.3)	11.3 (0.0-24.0)
C <sub>min</sub> (ng/dL)	361 ± 149	505 ± 233	235 ± 122
T <sub>min</sub> * (hr)	8.0 (0.0-24.1)	8.0 (0.0-25.8)	2.0 (0.0-24.2)
Fluc Index (ratio)	0.857 ± 0.331	0.895 ± 0.434	0.823 ± 0.289
Accum Ratio (ratio)	1.529 ± 0.726	1.911 ± 1.588	0.937 ± 0.354
<b>Day 90</b>	N = 65	N = 73	N = 64
C <sub>avg</sub> (ng/dL)	553 ± 247	792 ± 276	417 ± 157
C <sub>max</sub> (ng/dL)	846 ± 444	1204 ± 570	597 ± 242
T <sub>max</sub> * (hr)	4.0 (0.0-24.1)	7.9 (0.0-25.2)	8.1 (0.0-25.0)
C <sub>min</sub> (ng/dL)	354 ± 147	501 ± 193	213 ± 105
T <sub>min</sub> * (hr)	4.0 (0.0-25.3)	8.0 (0.0-24.8)	2.0 (0.0-24.0)
Fluc Index (ratio)	0.851 ± 0.402	0.859 ± 0.399	0.937 ± 0.442
Accum Ratio (ratio)	1.615 ± 0.859	1.927 ± 1.310	0.971 ± 0.453
<b>Day 180</b>	N = 63	N = 68	N = 45
C <sub>avg</sub> (ng/dL)	520 ± 227	722 ± 242	403 ± 163
C <sub>max</sub> (ng/dL)	779 ± 359	1091 ± 437	580 ± 240
T <sub>max</sub> * (hr)	4.0 (0.0-24.0)	7.9 (0.0-24.0)	10.0 (0.0-24.0)
C <sub>min</sub> (ng/dL)	348 ± 164	485 ± 184	223 ± 114
T <sub>min</sub> * (hr)	11.9 (0.0-24.0)	11.8 (0.0-27.4)	2.0 (0.0-25.7)
Fluc Index (ratio)	0.845 ± 0.379	0.829 ± 0.392	0.891 ± 0.319
Accum Ratio (ratio)	1.523 ± 1.024	1.897 ± 2.123	0.954 ± 0.4105

\*Median (Range)

**Table 8(f): Testosterone Pharmacokinetic Parameters on Days 30, 90, 180  
by Final Treatment Group (Mean  $\pm$  SD)**

	Doses Received During Initial $\Rightarrow$ Extended Treatment Phases				
	5.0 g/day T-gel	5.0 $\Rightarrow$ 7.5 g/day T-gel	10.0 $\Rightarrow$ 7.5 g/day T-gel	10.0 g/day T-gel	T-patch
<b>Day 30</b>	N = 47	N = 19	N = 19	N = 55	N = 70
C <sub>avg</sub> (ng/dL)	604 $\pm$ 288	472 $\pm$ 148	946 $\pm$ 399	739 $\pm$ 230	419 $\pm$ 163
C <sub>max</sub> (ng/dL)	941 $\pm$ 509	716 $\pm$ 294	1409 $\pm$ 556	1128 $\pm$ 436	576 $\pm$ 223
T <sub>max</sub> * (hr)	7.9 (0.0-24.0)	8.0 (0.0-24.0)	8.0 (0.0-24.3)	7.8 (0.0-24.3)	11.3 (0.0-24.0)
C <sub>min</sub> (ng/dL)	387 $\pm$ 159	296 $\pm$ 97	600 $\pm$ 339	471 $\pm$ 175	235 $\pm$ 122
T <sub>min</sub> * (hr)	8.1 (0.0-24.1)	1.7 (0.0-24.1)	11.4 (0.0-24.1)	8.0 (0.0-25.8)	2.0 (0.0-24.2)
Fluc Index (ratio)	0.861 $\pm$ 0.341	0.846 $\pm$ 0.315	0.927 $\pm$ 0.409	0.884 $\pm$ 0.445	0.823 $\pm$ 0.289
Accum Ratio (ratio)	1.543 $\pm$ 0.747	1.494 $\pm$ 0.691	2.053 $\pm$ 1.393	1.864 $\pm$ 1.657	0.937 $\pm$ 0.354
<b>Day 90</b>	N = 45	N = 20	N = 18	N = 55	N = 64
C <sub>avg</sub> (ng/dL)	596 $\pm$ 266	455 $\pm$ 164	859 $\pm$ 298	771 $\pm$ 268	417 $\pm$ 157
C <sub>max</sub> (ng/dL)	931 $\pm$ 455	654 $\pm$ 359	1398 $\pm$ 733	1141 $\pm$ 498	597 $\pm$ 242
T <sub>max</sub> * (hr)	3.8 (0.0-24.1)	7.7 (0.0-24.0)	7.9 (0.0-24.0)	7.9 (0.0-25.2)	8.1 (0.0-25.0)
C <sub>min</sub> (ng/dL)	384 $\pm$ 147	286 $\pm$ 125	532 $\pm$ 181	492 $\pm$ 197	213 $\pm$ 105
T <sub>min</sub> * (hr)	7.9 (0.0-25.3)	0.0 (0.0-24.0)	12.0 (0.0-24.1)	4.0 (0.0-24.8)	2.0 (0.0-24.0)
Fluc Index (ratio)	0.886 $\pm$ 0.391	0.771 $\pm$ 0.425	0.959 $\pm$ 0.490	0.826 $\pm$ 0.363	0.937 $\pm$ 0.442
Accum Ratio (ratio)	1.593 $\pm$ 0.813	1.737 $\pm$ 1.145	1.752 $\pm$ 0.700	1.952 $\pm$ 1.380	0.971 $\pm$ 0.453
<b>Day 180</b>	N = 44	N = 18	N = 19	N = 48	N = 41
C <sub>avg</sub> (ng/dL)	555 $\pm$ 225	450 $\pm$ 219	744 $\pm$ 320	713 $\pm$ 209	408 $\pm$ 165
C <sub>max</sub> (ng/dL)	803 $\pm$ 347	680 $\pm$ 369	1110 $\pm$ 468	1083 $\pm$ 434	578 $\pm$ 245
T <sub>max</sub> * (hr)	5.8 (0.0-24.0)	2.0 (0.0-24.0)	7.8 (0.0-24.0)	7.7 (0.0-24.0)	10.6 (0.0-24.0)
C <sub>min</sub> (ng/dL)	371 $\pm$ 165	302 $\pm$ 150	505 $\pm$ 233	485 $\pm$ 156	222 $\pm$ 116
T <sub>min</sub> * (hr)	11.9 (0.0-24.0)	9.9 (0.0-24.0)	12.0 (0.0-24.0)	8.0 (0.0-27.4)	2.0 (0.0-25.7)
Fluc Index (ratio)	0.853 $\pm$ 0.402	0.833 $\pm$ 0.335	0.824 $\pm$ 0.298	0.818 $\pm$ 0.421	0.866 $\pm$ 0.311
Accum Ratio (ratio)	1.541 $\pm$ 0.917	NA	NA	2.061 $\pm$ 2.445	0.969 $\pm$ 0.415

\*Median (range)

### (5) Dose Proportionality for AndroGel®

Table 8(g) shows the increase in AUC<sub>0-24</sub> on days 30, 90, and 180 from the pretreatment baseline (net AUC<sub>0-24</sub>). In order to assess dose-proportionality, the bioequivalence assessment was performed on the log-transformed AUCs using “treatment” as the only factor. The AUCs were compared after subtracting away the AUC contribution from the endogenous secretion of testosterone (the AUC on day 0) and adjusting for the two-fold difference in applied doses. The AUC ratio on day 30 was 0.95 (90% C.I.: 0.75-1.19) and on day 90 was 0.92 (90% C.I.: 0.73-1.17). When the day 30 and day 90 data was combined, the AUC ratio was 0.93 (90% C.I.: 0.79-1.10).

The data shows dose proportionality for AndroGel® treatment. The geometric mean for the increase in AUC<sub>0-24</sub> from day 0 to day 30 or day 90 was twice as great for the 10.0 g/day



group as for the 5.0 g/day group. A 125 ng/dL mean increase in serum testosterone  $C_{avg}$  level was produced by each 2.5 g/day of AndroGel<sup>®</sup>. In other words, the data shows that 0.1 g/day of AndroGel<sup>®</sup> produced, on the average, a 5 ng/dL increase in serum testosterone concentration. This dose proportionality aids dosing adjustment by the physician. Because AndroGel<sup>®</sup> is provided in 2.5 g packets (containing 25 mg of testosterone), each 2.5 g packet will produce, on average, a 125 ng/dL increase in the  $C_{avg}$  for serum total testosterone.

**Table 8(g): Net AUC<sub>0-24</sub> (nmol\*h/L) on Days 30, 90, and 180 after Transdermal Testosterone Application**

	<b>T Patch</b>	<b>T gel 5.0 g/day</b>	<b>T gel 10.0 g/day</b>
Day 30	154 ± 18	268 ± 28	446 ± 30
Day 90	157 ± 20	263 ± 29	461 ± 28
Day 180	160 ± 25	250 ± 32	401 ± 27

The increase in AUC<sub>0-24</sub> from pretreatment baseline achieved by the 10.0 g/day and the 5.0 g/day groups were approximately 2.7 and 1.7 fold higher than that resulting from application of the testosterone patch.

## **b. Pharmacokinetics of Serum Free Testosterone Concentration**

### **(1) Methods**

Serum free testosterone was measured by RIA of the dialysate, after an overnight equilibrium dialysis, using the same RIA reagents as the testosterone assay. The LLQ of serum free testosterone, using the equilibrium dialysis method, was estimated to be 22 pmol/L. When steroid free serum was spiked with increasing doses of testosterone in the adult male range, increasing amounts of free testosterone were recovered with a coefficient of variation that ranged from 11.0-18.5%. The intra- and interassay coefficients of free testosterone were 15% and 16.8% for adult normal male values, respectively. As estimated by the UCLA-Harbor Medical

Center, free testosterone concentrations range from 3.48-17.9 ng/dL (121-620 pmol/L) in normal adult men.

14111796V-2

## (2) Pharmacokinetic Results

In general, as shown in Table 9, the pharmacokinetic parameters of serum free testosterone mirrored that of serum total testosterone as described above. At baseline (day 0), the mean serum free testosterone concentrations ( $C_{avg}$ ) were similar in all three groups which were at the lower limit of the adult male range. The maximum serum free testosterone concentration occurred between 8 and 10 a.m., and the minimum about 8 to 16 hours later. This data is consistent with the mild diurnal variation of serum testosterone.

FIG. 6(a) shows the 24-hour pharmacokinetic profiles for the three treatment groups on day 1. After application of the testosterone patch, the serum free testosterone levels peaked at 12 hours about 4 hours earlier than those achieved by the AndroGel<sup>®</sup> groups. The serum free testosterone levels then declined in the testosterone patch group whereas in the AndroGel<sup>®</sup> groups, the serum free testosterone levels continued to rise.

FIGS. 6(b) and 6(c) show the pharmacokinetic profiles of free testosterone in the AndroGel<sup>®</sup>-treated groups resembled the unique testosterone profiles on days 30 and 90. After AndroGel<sup>®</sup> application, the mean serum free testosterone levels in the three groups were within normal range. Similar to the total testosterone results, the free testosterone  $C_{avg}$  achieved by the 10.0 g/day group was 1.4 fold higher than the 5.0 g/day group and 1.7 fold higher than the testosterone patch group. Moreover, the accumulation ratio for the testosterone patch was significantly less than that of the 5.0 g/day AndroGel<sup>®</sup> group and the 10.0 g/day AndroGel<sup>®</sup> group.

FIG. 6(d) shows the free testosterone concentrations by final treatment groups on day 180. In general, the free testosterone concentrations exhibited a similar pattern as serum testosterone. The 24-hour pharmacokinetic parameters were similar to those on days 30 and 90 in those subjects who remained in the three original randomized groups. Again, in the subjects

titrated to receive 7.5 g/day of AndroGel<sup>®</sup>, the group was not homogenous. The free testosterone  $C_{avg}$  in the patients with doses adjusted upwards from 5.0 to 7.5 g/day remained 29% lower than those of subjects remaining in the 5.0 g/day group. The free testosterone  $C_{avg}$  in the patients whose doses were decreased from 10.0 to 7.5 g/day was 11% higher than those in remaining in the 10.0 g/day group.

FIGS. 6(e)-(g) show the free testosterone concentrations in the three groups of subjects throughout the 180-day treatment period. Again, the free testosterone levels followed that of testosterone. The mean free testosterone levels in all three groups were within the normal range with the 10.0 g/day group maintaining higher free testosterone levels than both the 5.0 g/day and the testosterone patch groups.

**Table 9: Free Testosterone Pharmacokinetic Parameters by Final Treatment (Mean  $\pm$  SD)**

	Doses Received During Initial => Extended Treatment Phases				
	5.0 g/day T-gel	5.0 => 7.5 g/day T-gel	10.0 => 7.5 g/day T-gel	10/0 g/day T gel	T-patch
<b>Day 0</b>	N = 53	N = 20	N = 20	N = 58	N = 76
Cavg (ng/dL)	4.52 $\pm$ 3.35	4.27 $\pm$ 3.45	4.64 $\pm$ 3.10	4.20 $\pm$ 3.33	4.82 $\pm$ 3.64
Cmax (ng/dL)	5.98 $\pm$ 4.25	6.06 $\pm$ 5.05	6.91 $\pm$ 4.66	5.84 $\pm$ 4.36	6.57 $\pm$ 4.90
Tmax* (hr)	4.0 (0.0-24.5)	2.0 (0.0-24.0)	13.5 (0.0-24.2)	2.1 (0.0-24.1)	3.8 (0.0-24.0)
Cmin (ng/dL)	3.23 $\pm$ 2.74	3.10 $\pm$ 2.62	3.14 $\pm$ 2.14	3.12 $\pm$ 2.68	3.56 $\pm$ 2.88
Tmin* (hr)	8.0 (0.0-24.2)	9.9 (0.0-16.0)	4.0 (0.0-23.3)	8.0 (0.0-24.0)	7.9 (0.0-24.0)
Fluc Index (ratio)	0.604 $\pm$ 0.342	0.674 $\pm$ 0.512	0.756 $\pm$ 0.597	0.634 $\pm$ 0.420	0.614 $\pm$ 0.362
<b>Day 1</b>	N = 53	N = 20	N = 19	N = 57	N = 74
Cavg (ng/dL)	7.50 $\pm$ 4.83	6.80 $\pm$ 4.82	9.94 $\pm$ 5.04	8.93 $\pm$ 6.09	9.04 $\pm$ 4.81
Cmax (ng/dL)	10.86 $\pm$ 7.45	10.10 $\pm$ 7.79	15.36 $\pm$ 7.31	13.20 $\pm$ 8.61	12.02 $\pm$ 6.14
Tmax* (hr)	16.0 (0.0-25.3)	13.9 (0.0-24.3)	15.7 (2.0-24.0)	23.5 (1.8-24.3)	12.0 (1.8-24.0)
Cmin (ng/dL)	4.30 $\pm$ 3.33	3.69 $\pm$ 3.24	3.88 $\pm$ 2.73	4.40 $\pm$ 3.94	4.67 $\pm$ 3.52
Tmin* (hr)	0.0 (0.0-24.1)	1.8 (0.0-24.0)	0.0 (0.0-24.2)	0.0 (0.0-23.9)	0.0 (0.0-24.0)
<b>Day 30</b>	N = 47	N = 19	N = 19	N = 55	N = 70
Cavg (ng/dL)	11.12 $\pm$ 6.22	7.81 $\pm$ 3.94	16.18 $\pm$ 8.18	13.37 $\pm$ 7.13	8.12 $\pm$ 4.15
Cmax (ng/dL)	16.93 $\pm$ 10.47	11.62 $\pm$ 6.34	25.14 $\pm$ 10.80	19.36 $\pm$ 9.75	11.48 $\pm$ 5.78
Tmax* (hr)	8.0 (0.0-27.8)	8.0 (0.0-26.3)	8.0 (0.0-24.3)	8.0 (0.0-24.3)	8.0 (0.0-24.0)
Cmin (ng/dL)	6.99 $\pm$ 3.82	4.78 $\pm$ 3.10	9.99 $\pm$ 7.19	8.25 $\pm$ 5.22	4.31 $\pm$ 3.20
Tmin* (hr)	4.0 (0.0-24.1)	3.5 (0.0-24.1)	11.4 (0.0-24.1)	7.8 (0.0-25.8)	2.0 (0.0-24.8)
Fluc Index (ratio)	0.853 $\pm$ 0.331	0.872 $\pm$ 0.510	1.051 $\pm$ 0.449	0.861 $\pm$ 0.412	0.929 $\pm$ 0.311
Accum Ratio (ratio)	1.635 $\pm$ 0.820	1.479 $\pm$ 0.925	2.065 $\pm$ 1.523	1.953 $\pm$ 1.626	0.980 $\pm$ 0.387

	Doses Received During Initial => Extended Treatment Phases				
	5.0 g/day T-gel	5.0 => 7.5 g/day T-gel	10.0 => 7.5 g/day T-gel	10/0 g/day T gel	T-patch
<b>Day 90</b>	N = 45	N = 20	N = 18	N = 55	N = 64
Cavg (ng/dL)	12.12 ± 7.78	8.06 ± 3.78	17.65 ± 8.62	13.11 ± 5.97	8.50 ± 5.04
Cmax (ng/dL)	18.75 ± 12.90	10.76 ± 4.48	25.29 ± 12.42	18.61 ± 8.20	12.04 ± 6.81
Tmax* (hr)	4.0 (0.0-24.0)	9.7 (0.0-24.0)	8.0 (0.0-24.0)	8.0 (0.0-25.2)	11.6 (0.0-25.0)
Cmin (ng/dL)	7.65 ± 4.74	4.75 ± 2.86	10.56 ± 6.07	8.40 ± 4.57	4.38 ± 3.70
Tmin* (hr)	8.0 (0.0-24.0)	1.9 (0.0-24.0)	5.9 (0.0-24.1)	4.0 (0.0-24.8)	2.0 (0.0-24.1)
Fluc Index (ratio)	0.913 ± 0.492	0.815 ± 0.292	0.870 ± 0.401	0.812 ± 0.335	0.968 ± 0.402
Accum Ratio (ratio)	1.755 ± 0.983	1.916 ± 1.816	1.843 ± 0.742	2.075 ± 1.866	1.054 ± 0.498
<b>Day 180</b>	N = 44	N = 18	N = 19	N = 48	N = 41
Cavg (ng/dL)	11.01 ± 5.24	7.80 ± 4.63	14.14 ± 7.73	12.77 ± 5.70	7.25 ± 4.90
Cmax (ng/dL)	16.21 ± 7.32	11.36 ± 6.36	22.56 ± 12.62	18.58 ± 9.31	10.17 ± 5.90
Tmax* (hr)	7.9 (0.0-24.0)	2.0 (0.0-23.9)	7.8 (0.0-24.0)	8.0 (0.0-24.0)	11.1 (0.0-24.0)
Cmin (ng/dL)	7.18 ± 3.96	5.32 ± 4.06	9.54 ± 6.45	8.23 ± 4.01	3.90 ± 4.20
Tmin* (hr)	9.9 (0.0-24.2)	7.9 (0.0-24.0)	8.0 (0.0-23.2)	11.8 (0.0-27.4)	2.5 (0.0-25.7)
Fluc Index (ratio)	0.897 ± 0.502	0.838 ± 0.378	0.950 ± 0.501	0.815 ± 0.397	0.967 ± 0.370
Accum Ratio (ratio)	1.712 ± 1.071	NA	NA	2.134 ± 1.989	1.001 ± 0.580

\*Median (Range)

### c. Serum DHT Concentrations

Serum DHT was measured by RIA after potassium permanganate treatment of the sample followed by extraction. The methods and reagents of the DHT assay were provided by DSL (Webster, TX). The cross reactivities of the antiserum used in the RIA for DHT were 6.5% for 3- $\beta$ -androstanediol, 1.2% for 3- $\alpha$ -androstanediol, 0.4% for 3- $\alpha$ -androstanediol glucuronide, and 0.4% for testosterone (after potassium permanganate treatment and extraction), and less than 0.01% for other steroids tested. This low cross-reactivity against testosterone was further confirmed by spiking steroid free serum with 35 nmol/L (1,000 pg/dL) of testosterone and taking the samples through the DHT assay. The results even on spiking with over 35 nmol/L of testosterone was measured as less than 0.1 nmol/L of DHT. The LLQ of serum DHT in the assay was 0.43 nmol/L. The mean accuracy (recovery) of the DHT assay determined by spiking steroid free serum with varying amounts of DHT from 0.43 nmol/L to 9 nmol/L was 101% and ranged from 83 to 114%. The intra-assay and inter-assay coefficients of variation for the DHT

assay were 7.8 and 16.6%, respectively, for the normal adult male range. The normal adult male range of DHT was 30.7-193.2 ng/dL (1.06 to 6.66 nmol/L ) as determined by the UCLA-Harbor Medical Center.

As shown in Table 10, the pretreatment mean serum DHT concentrations were between 36 and 42 ng/dL, which were near the lower limit of the normal range in all three initial treatment groups. None of the patients had DHT concentrations above the upper limit of the normal range on the pretreatment day, although almost half (103 patients) had concentrations less than the lower limit.

FIG. 7 shows that after treatment, the differences between the mean DHT concentrations associated with the different treatment groups were statistically significant, with patients receiving AndroGel® having a higher mean DHT concentration than the patients using the patch and showing dose-dependence in the mean serum DHT concentrations. Specifically, after testosterone patch application mean serum DHT levels rose to about 1.3 fold above the baseline. In contrast, serum DHT increased to 3.6 and 4.8 fold above baseline after application of 5.0 g/day and 10.0 g/day of AndroGel®, respectively.

**Table 10: DHT Concentrations (ng/dL)  
on Each of the Observation Days  
By Initial Treatment (Mean ± SD)**

	Day 0	Day 30	Day 60	Day 90	Day 120	Day 150	Day 180
<b>5.0 g/day T-gel</b>	N = 73 36.0 ± 19.9	N = 69 117.6 ± 74.9	N = 70 122.4 ± 99.4	N = 67 130.1 ± 99.2	N = 65 121.8 ± 89.2	N = 63 144.7 ± 110.5	N = 65 143.7 ± 105.9
<b>10.0 g/day T-gel</b>	N = 78 42.0 ± 29.4	N = 78 200.4 ± 127.8	N = 74 222.0 ± 126.6	N = 75 207.7 ± 111.0	N = 68 187.3 ± 97.3	N = 67 189.1 ± 102.4	N = 71 206.1 ± 105.9
<b>T-Patch</b>	N = 76 37.4 ± 21.4	N = 73 50.8 ± 34.6	N = 68 49.3 ± 27.2	N = 66 43.6 ± 26.9	N = 49 53.0 ± 52.8	N = 46 54.0 ± 42.5	N = 49 52.1 ± 34.3
Across RX	0.6041	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001

The increase in DHT concentrations are likely attributed to the concentration and location of 5α-reductase in the skin. For example, the large amounts of 5α-reductase in the scrotal skin presumably causes an increase in DHT concentrations in the TESTODERM® patch. In contrast,

the ANDRODERM<sup>®</sup> and TESTODERM TTS<sup>®</sup> patches create little change in DHT levels because the surface area of the patch is small and little 5 $\alpha$ -reductase is located in nonscrotal skin. AndroGel<sup>®</sup> presumably causes an increase in DHT levels because the gel is applied to a relatively large skin area and thus exposes testosterone to greater amounts of the enzyme.

5 To date, elevated DHT levels have not been reported to have any adverse clinical effects. Moreover, there is some evidence to suggest that increased DHT levels may inhibit prostate cancer.

#### d. DHT/T Ratio

The UCLA-Harbor Medical Center reports a DHT/T ratio of 0.052-0.328 for normal adult men. In this example, the mean ratios for all three treatments were within the normal range on day 0. As shown in FIG. 8 and Table 11, there were treatment and concentration-dependent increases observed over the 180-day period. Specifically, the AndroGel<sup>®</sup> treatment groups showed the largest increase in DHT/T ratio. However, the mean ratios for all of the treatment groups remained within the normal range on all observation days.

**Table 11: DHT/T Ratio  
on Each of the Observation Days  
By Initial Treatment (Mean  $\pm$  SD)**

	Day 0	Day 30	Day 60	Day 90	Day 120	Day 150	Day 180
<b>5.0 g/day T-gel</b>	N = 73 0.198 $\pm$ 0.137	N = 68 0.230 $\pm$ 0.104	N = 70 0.256 $\pm$ 0.132	N = 67 0.248 $\pm$ 0.121	N = 65 0.266 $\pm$ 0.119	N = 62 0.290 $\pm$ 0.145	N = 64 0.273 $\pm$ 0.160
<b>10.0 g/day T-gel</b>	N = 78 0.206 $\pm$ 0.163	N = 77 0.266 $\pm$ 0.124	N = 74 0.313 $\pm$ 0.160	N = 74 0.300 $\pm$ 0.131	N = 68 0.308 $\pm$ 0.145	N = 67 0.325 $\pm$ 0.142	N = 71 0.291 $\pm$ 0.124
<b>T-Patch</b>	N = 76 0.204 $\pm$ 0.135	N = 73 0.192 $\pm$ 0.182	N = 68 0.175 $\pm$ 0.102	N = 65 0.175 $\pm$ 0.092	N = 49 0.186 $\pm$ 0.134	N = 46 0.223 $\pm$ 0.147	N = 46 0.212 $\pm$ 0.160
<b>Across RX</b>	0.7922	0.0001	0.0001	0.0001	0.0001	0.0001	0.0002

#### e. Total Androgen (DHT + T)

20 The UCLA-Harbor Medical Center has determined that the normal total androgen concentration is 372 to 1,350 ng/dL. As shown in FIG. 9 and Table 12, the mean pre-dose total androgen concentrations for all three treatments were below the lower limit of the normal range

on pretreatment day 0. The total androgen concentrations for both AndroGel<sup>®</sup> groups were within the normal range on all treatment observation days. In contrast, the mean concentrations for patients receiving the testosterone patch was barely within the normal range on day 60 and 120, but were below the lower normal limit on days 30, 90, 150, and 180.

5 **Table 12: Total Androgen (DHT +T) (ng/dL)  
on Each of the Observation Days  
By Initial Treatment (Mean ± SD)**

	Day 0	Day 30	Day 60	Day 90	Day 120	Day 150	Day 180
<b>5.0 g/day T-gel</b>	N = 73 281 ± 150	N = 68 659 ± 398	N = 70 617 ± 429	N = 67 690 ± 431	N = 65 574 ± 331	N = 62 631 ± 384	N = 64 694 ± 412
<b>10.0 g/day T-gel</b>	N = 78 307 ± 180	N = 77 974 ± 532	N = 74 1052 ± 806	N = 74 921 ± 420	N = 68 827 ± 361	N = 67 805 ± 383	N = 71 944 ± 432
<b>T-Patch</b>	N = 76 282 ± 159	N = 73 369 ± 206	N = 68 392 ± 229	N = 65 330 ± 173	N = 49 378 ± 250	N = 46 364 ± 220	N = 46 355 ± 202
<b>Across RX</b>	0.7395	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001

#### f. E<sub>2</sub> Concentrations

Serum E<sub>2</sub> levels were measured by a direct assay without extraction with reagents from ICN (Costa Mesa, CA). The intra-assay and inter-assay coefficients of variation of E<sub>2</sub> were 6.5 and 7.1% respectively. The UCLA-Harbor Medical Center reported an average E<sub>2</sub> concentration ranging from 7.1 to 46.1 pg/mL (63 to 169 pmol/L) for normal adult male range. The LLQ of the E<sub>2</sub> was 18 pmol/L. The cross reactivities of the E<sub>2</sub> antibody were 6.9% for estrone, 0.4% for equilinenin, and less than 0.01% for all other steroids tested. The accuracy of the E<sub>2</sub> assay was assessed by spiking steroid free serum with increasing amount of E<sub>2</sub> (18 to 275 pmol/L). The mean recovery of E<sub>2</sub> compared to the amount added was 99.1% and ranged from 95 to 101%.

FIG. 10 depicts the E<sub>2</sub> concentrations throughout the 180-day study. The pretreatment mean E<sub>2</sub> concentrations for all three treatment groups were 23-24 pg/mL. During the study, the E<sub>2</sub> levels increased by an average 9.2% in the testosterone patch during the treatment period, 30.9% in the 5.0 g/day AndroGel<sup>®</sup> group, and 45.5% in the 10.0 g/day AndroGel<sup>®</sup> group. All of the mean concentrations fell within the normal range.



**Table 13: Estradiol Concentration (pg/mL)  
on Each of the Observation Days  
By Initial Treatment (Mean ± SD)**

	Day 0	Day 30	Day 60	Day 90	Day 120	Day 150	Day 180
5.0 g/day T-gel	N = 73 23.0 ± 9.2	N = 69 29.2 ± 11.0	N = 68 28.1 ± 10.0	N = 67 31.4 ± 11.9	N = 64 28.8 ± 9.9	N = 65 30.8 ± 12.5	N = 65 32.3 ± 13.8
10.0 g/day T-gel	N = 78 24.5 ± 9.5	N = 78 33.7 ± 11.5	N = 74 36.5 ± 13.5	N = 75 37.8 ± 13.3	N = 71 34.6 ± 10.4	N = 66 35.0 ± 11.1	N = 71 36.3 ± 13.9
T-Patch	N = 76 23.8 ± 8.2	N = 72 25.8 ± 9.8	N = 68 24.8 ± 8.0	N = 66 25.7 ± 9.8	N = 50 25.7 ± 9.4	N = 49 27.0 ± 9.2	N = 49 26.9 ± 9.5
Across RX	0.6259	0.0001	0.0001	0.0001	0.0001	0.0009	0.0006

5 E<sub>2</sub> is believed to be important for the maintenance of normal bone. In addition, E<sub>2</sub> has a positive effect on serum lipid profiles.

#### g. Serum SHBG Concentrations

Serum SHBG levels were measured with a fluoroimmunoassay ("FIA") obtained from Delfia (Wallac, Gaithersburg, MD). The intra- and interassay coefficients were 5% and 12% respectively. The LLQ was 0.5 nmol/L. The UCLA-Harbor Medical Center determined that the adult normal male range for the SHBG assay is 0.8 to 46.6 nmol/L.

As shown in FIG. 11 and Table 11, the serum SHBG levels were similar and within the normal adult male range in the three treatment groups at baseline. None of the treatment groups showed major changes from these the baseline on any of the treatment visit days. After testosterone replacement serum SHBG levels showed a small decrease in all three groups. The most marked change occurred in the 10.0 g/day AndroGel<sup>®</sup> group.

**Table 14: SHBG Concentration (nmol/L)  
on Each of the Observation Days  
By Initial Treatment (Mean ± SD)**

	Day 0	Day 30	Day 60	Day 90	Day 120	Day 150	Day 180
5.0 g/day T-gel	N = 73 26.2 ± 14.9	N = 69 24.9 ± 14.0	N = 69 25.9 ± 14.4	N = 67 25.5 ± 14.7	N = 66 25.2 ± 14.1	N = 65 24.9 ± 12.9	N = 65 24.2 ± 13.6
10.0 g/day T-gel	N = 78 26.6 ± 17.8	N = 78 24.8 ± 14.5	N = 75 25.2 ± 15.5	N = 75 23.6 ± 14.7	N = 72 25.5 ± 16.5	N = 68 23.8 ± 12.5	N = 71 24.0 ± 14.5
T-Patch	N = 76 30.2 ± 22.6	N = 72 28.4 ± 21.3	N = 68 28.2 ± 23.8	N = 66 28.0 ± 23.6	N = 50 26.7 ± 16.0	N = 49 26.7 ± 16.4	N = 49 25.8 ± 15.1
Across RX	0.3565	0.3434	0.5933	0.3459	0.8578	0.5280	0.7668

## h. Gonadotropins

Serum FSH and LH were measured by highly sensitive and specific solid-phase FIA assays with reagents provided by Delfia (Wallac, Gaithersburg, MD). The intra-assay coefficient of variations for LH and FSH fluoroimmunoassays were 4.3 and 5.2%, respectively; and the interassay variations for LH and FSH were 11.0% and 12.0%, respectively. For both LH and FSH assays, the LLQ was determined to be 0.2 IU/L. All samples obtained from the same subject were measured in the same assay. The UCLA-Harbor Medical Center reports that the adult normal male range for LH is 1.0-8.1 U/L and for FSH is 1.0-6.9U/L.

### (1) FSH

Table 15(a)-(d) shows the concentrations of FSH throughout the 180-day treatment depending on the cause of hypogonadism: (1) primary, (2) secondary, (3) age-associated, or (4) unknown.

As discussed above, patients with primary hypogonadism have an intact feedback inhibition pathway, but the testes do not secrete testosterone. As a result, increasing serum testosterone levels should lead to a decrease in the serum FSH concentrations. In this example, a total of 94 patients were identified as having primary hypogonadism. For these patients, the mean FSH concentrations in the three treatment groups on day 0 were 21-26 mIU/mL, above the upper limit of the normal range. As shown in FIG. 12(a) and Table 15(a), the mean FSH concentrations decreased during treatment in all three treatment regimens. However, only the 10.0 g/day AndroGel® group reduced the mean concentrations to within the normal range during the first 90 days of treatment. Treatment with the 10.0 g/day AndroGel® group required approximately 120 days to reach steady state. The mean FSH concentration in patients applying 5.0 g/day of AndroGel® showed an initial decline that was completed by day 30 and another declining phase at day 120 and continuing until the end of treatment. Mean FSH concentrations

in the patients receiving the testosterone patch appeared to reached steady state after 30 days but were significantly higher than the normal range.

**Table 15(a): FSH Concentrations (mIU/mL) on Each of the Observation Days by Initial Treatment Group for Patients Having Primary Hypogonadism (Mean  $\pm$  SD)**

	N	5 g/day	N	10 g/day	N	T-patch
Day 0	26	21.6 $\pm$ 21.0	33	20.9 $\pm$ 15.9	34	25.5 $\pm$ 25.5
Day 30	23	10.6 $\pm$ 15.0	34	10.6 $\pm$ 14.1	31	21.4 $\pm$ 24.6
Day 60	24	10.8 $\pm$ 16.9	32	7.2 $\pm$ 12.6	31	21.7 $\pm$ 23.4
Day 90	24	10.4 $\pm$ 19.7	31	5.7 $\pm$ 10.1	30	19.5 $\pm$ 20.0
Day 120	24	8.1 $\pm$ 15.2	28	4.6 $\pm$ 10.2	21	25.3 $\pm$ 28.4
Day 150	22	6.7 $\pm$ 15.0	29	5.3 $\pm$ 11.0	21	18.6 $\pm$ 24.0
Day 180	24	6.2 $\pm$ 11.3	28	5.3 $\pm$ 11.2	22	24.5 $\pm$ 27.4

Patients with secondary hypogonadism have a deficient testosterone negative feedback system. As shown in FIG. 12(b), of 44 patients identified as having secondary hypogonadism, the mean FSH concentrations decreased during treatment, although the decrease over time was not statistically significant for the testosterone patch. The patients in the 5.0 g/day AndroGel<sup>®</sup> group showed a decrease in the mean FSH concentration by about 35% by day 30, with no further decrease evident by day 60. Beyond day 90, the mean FSH concentration in the patients appeared to slowly return toward the pretreatment value. By day 30, all of the 10.0 g/day AndroGel<sup>®</sup> group had FSH concentrations less than the lower limit.

**Table 15(b): FSH Concentrations (mIU/mL) on Each of the Observation Days by Initial Treatment Group for Patients Having Secondary Hypogonadism (Mean  $\pm$  SD)**

	N	5 g/day	N	10 g/day	N	T-patch
Day 0	17	4.2 $\pm$ 6.6	12	2.1 $\pm$ 1.9	15	5.1 $\pm$ 9.0
Day 30	16	2.8 $\pm$ 5.9	12	0.2 $\pm$ 0.1	14	4.2 $\pm$ 8.0
Day 60	17	2.8 $\pm$ 6.1	12	0.2 $\pm$ 0.1	13	4.2 $\pm$ 7.4
Day 90	15	2.9 $\pm$ 5.6	12	0.2 $\pm$ 0.1	14	4.9 $\pm$ 9.0
Day 120	14	3.0 $\pm$ 6.1	12	0.1 $\pm$ 0.1	12	6.1 $\pm$ 10.7
Day 150	14	3.5 $\pm$ 7.5	12	0.2 $\pm$ 0.2	11	4.6 $\pm$ 6.5
Day 180	14	3.7 $\pm$ 8.6	12	0.1 $\pm$ 0.1	12	4.9 $\pm$ 7.4

Twenty-five patients were diagnosed with age-associated hypogonadism. As shown in FIG. 12(c), the 5.0 g/day AndroGel® group had a mean pretreatment FSH concentration above the normal range. The mean concentration for this group was within the normal range by day 30 and had decreased more than 50% on days 90 and 180. The decrease in FSH mean concentration in the 10.0 g/day AndroGel® group showed a more rapid response. The concentrations in all six patients decreased to below the lower normal limit by day 30 and remained there for the duration of the study. The six patients who received the testosterone patch exhibited no consistent pattern in the mean FSH level; however, there was an overall trend towards lower FHS levels with continued treatment.

**Table 15(c): FSH Concentrations (mIU/mL) on Each of the Observation Days by Initial Treatment Group for Patients Having Age-Related Hypogonadism (Mean  $\pm$  SD)**

	N	5 g/day	N	10 g/day	N	T-patch
Day 0	13	8.0 ± 9.1	6	5.2 ± 1.9	6	4.7 ± 1.7
Day 30	12	4.6 ± 7.4	6	0.4 ± 0.3	6	3.7 ± 2.0
Day 60	12	3.9 ± 6.6	6	0.3 ± 0.3	4	4.3 ± 3.3
Day 90	11	3.8 ± 7.0	6	0.4 ± 0.7	4	3.5 ± 1.9
Day 120	11	4.2 ± 8.3	6	0.4 ± 0.7	4	4.2 ± 3.3
Day 150	11	4.3 ± 8.1	5	0.2 ± 0.2	4	3.4 ± 2.7
Day 180	11	4.0 ± 7.2	6	0.2 ± 0.2	4	2.7 ± 2.1

Sixty-four patients in the study suffered from unclassified hypogonadism. As shown in FIG. 12(d), the patients showed a marked and comparatively rapid FSH concentration decrease in all three groups, with the greatest decrease being in the 10.0 g/day AndroGel® group. The 10.0 g/day AndroGel® group produced nearly a 90% decrease in the mean FSH concentration by day 30 and maintained the effect to day 180. The 5.0 g/day AndroGel® group produced about a 75% drop in mean FSH concentration by day 30 and stayed at that level for the remainder of treatment. The 21 patients receiving the testosterone patch had a 50% decrease in the mean FSH

concentration by day 30, a trend that continued to day 90 when the concentration was about one-third of its pretreatment value.

**Table 15(d): Concentrations (mIU/mL) for FSH on Each of the Observation Days by Initial Treatment Group for Patients Having Unknown-Related Hypogonadism (Mean  $\pm$  SD)**

	N	5 g/day	N	10 g/day	N	T-patch
Day 0	17	4.0 $\pm$ 1.8	26	4.1 $\pm$ 1.6	21	3.7 $\pm$ 1.4
Day 30	17	1.1 $\pm$ 1.0	26	0.5 $\pm$ 0.5	21	1.8 $\pm$ 0.8
Day 60	16	1.1 $\pm$ 1.1	26	0.3 $\pm$ 0.3	18	1.6 $\pm$ 1.0
Day 90	17	1.1 $\pm$ 1.1	25	0.4 $\pm$ 0.7	18	1.2 $\pm$ 0.9
Day 120	16	1.2 $\pm$ 1.4	26	0.4 $\pm$ 0.6	12	1.4 $\pm$ 1.0
Day 150	17	1.4 $\pm$ 1.4	23	0.3 $\pm$ 0.5	13	1.4 $\pm$ 1.2
Day 180	16	1.0 $\pm$ 0.9	24	0.4 $\pm$ 0.4	11	1.3 $\pm$ 0.9

This data shows that feedback inhibition of FSH secretion functioned to some extent in all four subpopulations. The primary hypogonadal population showed a dose-dependency in both the extent and rate of the decline in FSH levels. The sensitivity of the feedback process appeared to be reduced in the secondary and age-associated groups in that only the highest testosterone doses had a significant and prolonged impact on FSH secretion. In contrast, the feedback inhibition pathway in the patients in the unclassified group was quite responsive at even the lowest dose of exogenous testosterone.

## (2) LH

The response of LH to testosterone was also examined separately for the same four subpopulations. Tables 16(a)-(d) shows the LH concentrations throughout the treatment period.

As shown in FIG. 13(a) and Table 16(a), the LH concentrations prior to treatment were about 175% of the upper limit of the normal range in primary hypogonadal patients. The mean LH concentrations decreased during treatment in all groups. However, only the AndroGel® groups decreased the mean LH concentrations enough to fall within the normal range. As with

FSH, the primary hypogonadal men receiving AndroGel® showed dose-dependence in both the rate and extent of the LH response.

**Table 16(a): Concentrations for LH (mIU/mL) on Each of the Observation Days for Patients Having Primary Hypogonadism (Summary of Mean  $\pm$  SD)**

	N	5 g/day	N	10 g/day	N	T-patch
Day 0	26	12.2 $\pm$ 12.1	33	13.9 $\pm$ 14.9	33	13.3 $\pm$ 14.3
Day 30	23	5.6 $\pm$ 7.6	34	5.9 $\pm$ 8.1	31	10.9 $\pm$ 12.9
Day 60	24	6.8 $\pm$ 9.0	32	4.8 $\pm$ 10.0	31	10.8 $\pm$ 11.8
Day 90	24	5.9 $\pm$ 9.5	31	4.2 $\pm$ 11.0	30	10.0 $\pm$ 11.7
Day 120	24	6.4 $\pm$ 11.9	28	3.8 $\pm$ 10.4	21	11.5 $\pm$ 11.5
Day 150	22	4.4 $\pm$ 8.5	29	4.0 $\pm$ 11.3	21	7.4 $\pm$ 6.0
Day 180	24	4.8 $\pm$ 6.8	28	4.0 $\pm$ 11.9	22	11.2 $\pm$ 10.5

The secondary hypogonadal men were less sensitive to exogenous testosterone. For the 44 patients identified as having secondary hypogonadism, the pretreatment mean concentrations were all within the lower limit normal range. The mean LH concentrations decreased during treatment with all three regimens as shown in FIG. 13(b) and Table 16(b).

**Table 16(b): Concentrations for LH (mIU/mL) on Each of the Observation Days for Patients Having Secondary Hypogonadism (Summary of Mean  $\pm$  SD)**

	N	5 g/day	N	10 g/day	N	T-patch
Day 0	17	1.8 $\pm$ 2.6	12	1.4 $\pm$ 1.8	15	1.6 $\pm$ 3.1
Day 30	16	1.1 $\pm$ 2.2	12	0.2 $\pm$ 0.2	14	0.4 $\pm$ 0.4
Day 60	17	1.4 $\pm$ 3.8	12	0.2 $\pm$ 0.2	13	0.6 $\pm$ 0.5
Day 90	15	1.2 $\pm$ 2.4	12	0.2 $\pm$ 0.2	14	0.7 $\pm$ 1.0
Day 120	14	1.6 $\pm$ 4.0	12	0.2 $\pm$ 0.2	12	0.8 $\pm$ 0.8
Day 150	14	1.6 $\pm$ 3.5	12	0.2 $\pm$ 0.2	11	1.2 $\pm$ 2.0
Day 180	14	1.5 $\pm$ 3.7	12	0.2 $\pm$ 0.2	12	1.4 $\pm$ 2.1

None of the 25 patients suffering from age-associated hypogonadism had pretreatment LH concentrations outside of the normal range as shown in FIG. 13(c) and Table 16(c). The overall time and treatment effects were significant for the AndroGel® patients but not those patients using the testosterone patch.

**Table 16(c): Concentrations for LH (mIU/mL) on Each of the Observation Days for Patients Having Age-Related Hypogonadism (Summary of Mean  $\pm$  SD)**

	N	5 g/day	N	10 g/day	N	T-patch
Day 0	13	3.2 $\pm$ 1.1	6	2.4 $\pm$ 1.8	6	2.9 $\pm$ 0.6
Day 30	12	1.1 $\pm$ 1.0	6	0.1 $\pm$ 0.0	6	1.8 $\pm$ 1.1
Day 60	12	0.8 $\pm$ 0.7	6	0.2 $\pm$ 0.3	5	3.4 $\pm$ 2.8
Day 90	11	0.9 $\pm$ 1.2	6	0.1 $\pm$ 0.0	4	2.3 $\pm$ 1.4
Day 120	11	1.0 $\pm$ 1.4	6	0.1 $\pm$ 0.0	4	2.2 $\pm$ 1.4
Day 150	11	1.3 $\pm$ 1.5	5	0.1 $\pm$ 0.0	4	1.9 $\pm$ 1.2
Day 180	11	1.8 $\pm$ 2.1	6	0.1 $\pm$ 0.0	4	1.4 $\pm$ 1.0

5 Of the 64 patients suffering from an unclassified hypogonadism, none of the patients had a pretreatment LH concentration above the upper limit. Fifteen percent, however, had pretreatment concentrations below the normal limit. The unclassified patients showed comparatively rapid LH concentration decreases in all treatment groups as shown in FIG. 13(d) and Table 16(d).

**Table 16(d): Concentrations for LH (mIU/mL) on Each of the Observation Days for Patients Having Unknown-Related Hypogonadism (Summary of Mean  $\pm$  SD)**

	N	5 g/day	N	10 g/day	N	T-patch
Day 0	17	1.8 $\pm$ 1.2	26	2.5 $\pm$ 1.5	21	2.5 $\pm$ 1.5
Day 30	17	0.3 $\pm$ 0.3	26	0.3 $\pm$ 0.3	21	1.3 $\pm$ 1.3
Day 60	17	0.4 $\pm$ 0.5	26	0.3 $\pm$ 0.3	18	1.2 $\pm$ 1.4
Day 90	17	0.5 $\pm$ 0.5	26	0.3 $\pm$ 0.4	18	1.0 $\pm$ 1.4
Day 120	17	0.4 $\pm$ 0.4	26	0.4 $\pm$ 0.5	12	1.2 $\pm$ 1.1
Day 150	17	0.8 $\pm$ 1.1	23	0.3 $\pm$ 0.4	13	1.1 $\pm$ 1.1
Day 180	15	0.3 $\pm$ 0.4	25	0.4 $\pm$ 0.4	11	1.5 $\pm$ 1.3

### (3) Summary: LH and FSH

15 Patients receiving AndroGel® or the testosterone patch achieve “hormonal steady state” only after long-term treatment. Specifically, data involving FSH and LH show that these hormones do not achieve steady-state until many weeks after treatment. Because testosterone concentrations are negatively inhibited by FSH and LG, testosterone levels do not achieve true steady state until these other hormones also achieve steady state. However, because these

hormones regulate only endogenous testosterone (which is small to begin with in hypogonadal men) in an intact feedback mechanism (which may not be present depending on the cause of hypogonadism), the level of FSH and/or LH may have little effect on the actual testosterone levels achieved. The net result is that the patients do not achieve a “hormonal steady state” for testosterone even though the  $C_{avg}$ ,  $C_{min}$ , and  $C_{max}$  for testosterone remains relative constant after a few days of treatment.

## **2. Bone Mineral Density (“BMD”) and Similar Markers**

### **a. BMD**

BMD was assessed by dual energy X-ray absorptiometry (“DEXA”) using Hologic QDR 2000 or 4500 A (Hologic, Waltham, MA) on days 0 and 180 in the lumbar spine and left hip regions. BMD of spine was calculated as the average of BMD at L1 to L4. BMD of the left hip, which included Ward’s triangle, was calculated by the average of BMD from neck, trochanter, and intertrochanter regions. The scans were centrally analyzed and processed at Hologic. BMD assessments were performed at 13 out of the 16 centers (206 out of 227 subjects) because of the lack of the specific DEXA equipment at certain sites.

Table 17 and FIGS. 14(a)-14(b) show that before treatment, the BMD of the hip or the spine was not different among the three treatment groups. Significant increases in BMD occurred only in subjects in the AndroGel® 10.0 g/day group and those who switched from AndroGel® 10.0 to 7.5 g/day groups. The increases in BMD were about 1% in the hip and 2% in the spine during the six-month period. Average increases in BMD of 0.6% and 1% in the hip and spine were seen in those receiving 5.0 g/day of AndroGel® but no increase was observed in the testosterone patch group.



**Table 17: BMD Concentrations on Day 0 and Day 180  
by Final Treatment Group Mean ( $\pm$  SD)**

Final Treatment Group	N	Day 0	N	Day 180	N	% Change from Day 0 to Day 180
<b>Hip</b>						
5.0 g/day T-gel	50	1.026 $\pm$ 0.145	41	1.022 $\pm$ 0.145	41	0.7 $\pm$ 2.1
5.0 to 7.5 g/day T-gel	16	1.007 $\pm$ 0.233	15	1.011 $\pm$ 0.226	15	1.0 $\pm$ 4.9
10.0 to 7.5 g/day T-gel	20	1.002 $\pm$ 0.135	19	1.026 $\pm$ 0.131	19	1.3 $\pm$ 2.4
10.0 g/day T-gel	53	0.991 $\pm$ 0.115	44	0.995 $\pm$ 0.130	44	1.1 $\pm$ 1.9
T-Patch	67	0.982 $\pm$ 0.166	37	0.992 $\pm$ 0.149	37	-0.2 $\pm$ 2.9
<b>Spine</b>						
5.0 g/day T-gel	50	1.066 $\pm$ 0.203	41	1.072 $\pm$ 0.212	41	1.0 $\pm$ 2.9
5.0 to 7.5 g/day T-gel	16	1.060 $\pm$ 0.229	15	1.077 $\pm$ 0.217	15	0.4 $\pm$ 5.5
10.0 to 7.5 g/day T-gel	19	1.049 $\pm$ 0.175	19	1.067 $\pm$ 0.175	18	1.4 $\pm$ 3.2
10.0 g/day T-gel	53	1.037 $\pm$ 0.126	44	1.044 $\pm$ 0.124	44	2.2 $\pm$ 3.1
T-Patch	67	1.058 $\pm$ 0.199	36	1.064 $\pm$ 0.205	36	-0.2 $\pm$ 3.4

Note: Day 0 and Day 180 are arithmetic means, while percent change is a geometric mean.

The baseline hip and spine BMD and the change in BMD on day 180 were not significantly correlated with the average serum testosterone concentration on day 0. The changes in BMD in the hip or spine after testosterone replacement were not significantly different in subjects with hypogonadism due to primary, secondary, aging, or unclassified causes; nor were they different between naive and previously testosterone replaced subjects. The change in BMD in the spine was negatively correlated with baseline BMD values, indicating that the greatest increase in BMD occurred in subjects with the lowest initial BMD. The increase in BMD in the hip (but not in the spine) after testosterone treatment was correlated with the change in serum testosterone levels.

#### **b. Bone Osteoblastic Activity Markers**

The results described above are supported by measurements of a number of serum and urine markers of bone formation. Specifically, the mean concentrations of the serum markers (PTH, SALP, osteocalcin, type I procollagen) generally increase during treatment in all treatment groups. In addition, the ratios of two urine markers of bone formation (N-telopeptide /creatinine ratio and calcium/creatinine ratio) suggests a decrease in bone resorption.

## (1) PTH (Parathyroid or Calciotropic Hormone)

Serum intact PTH was measured by two site immunoradiometric assay ("IRMA") kits from Nichol's Institute (San Juan Capistrano, CA). The LLC for the PTH assay was 12.5 ng/L. The intra- and inter-assay coefficients of variation were 6.9 and 9.6%, respectively. The UCLA-Harbor Medical Center has reported previously that the normal male adult range of PTH is 6.8 to 66.4 ng/L.

Table 18 provides the PTH concentrations over the 180-day study. FIG. 15 shows that the mean serum PTH levels were within the normal male range in all treatment groups at baseline. Statistically significant increases in serum PTH were observed in all subjects as a group at day 90 without inter-group differences. These increases in serum PTH were maintained at day 180 in all three groups.

**Table 18: PTH Concentrations on Each of the Observation Days  
by Final Treatment Group (Mean  $\pm$  SD)**

	N	5 g/day T-gel	N	5 $\Rightarrow$ 7.5 g/day T-gel	N	10 $\Rightarrow$ 7.5 g/day T-gel	N	10 g/day T-gel	N	T-Patch
Day 0	53	16.31 $\pm$ 8.81	20	17.70 $\pm$ 9.66	20	18.02 $\pm$ 8.18	58	14.99 $\pm$ 6.11	75	15.60 $\pm$ 6.57
Day 30	49	17.91 $\pm$ 10.36	20	18.33 $\pm$ 8.02	20	17.45 $\pm$ 5.67	58	18.04 $\pm$ 8.95	72	18.33 $\pm$ 10.92
Day 90	47	21.32 $\pm$ 11.47	20	21.25 $\pm$ 10.96	19	17.10 $\pm$ 6.04	54	20.01 $\pm$ 9.77	66	21.45 $\pm$ 13.71
Day 120	46	21.19 $\pm$ 11.42	19	21.42 $\pm$ 13.20	20	19.62 $\pm$ 9.96	50	22.93 $\pm$ 12.57	46	21.07 $\pm$ 11.44
Day 180	46	22.85 $\pm$ 12.89	19	21.34 $\pm$ 11.08	19	21.02 $\pm$ 10.66	51	25.57 $\pm$ 15.59	46	25.45 $\pm$ 16.54

## (2) SALP

SALP was quantitated by IRMA using reagents supplied by Hybritech (San Diego, CA). The LLQ for the SALP assay was 3.8  $\mu$ g/L.; and the intra- and inter-assay precision coefficients were 2.9 and 6.5%, respectively. The UCLA-Harbor Medical Center reported that the adult normal male concentration of SALP ranges from 2.4 to 16.6  $\mu$ g/L.

The pretreatment SALP concentrations were within the normal range. FIG. 16 and Table 19 show that SALP levels increased with testosterone treatment in the first 90 days and

reached statistical difference in the testosterone patch group. Thereafter serum SALP plateaued in all treatment groups.

**Table 19: SALP Concentrations on Each of the Observation Days  
by Final Treatment Group (Mean  $\pm$  SD)**

	N	5 g/day T-gel	N	5 $\Rightarrow$ 7.5 g/day T-gel	N	10 $\Rightarrow$ 7.5 g/day T-gel	N	10 g/day T-gel	N	T-Patch
Day 0	53	9.96 $\pm$ 5.61	20	12.36 $\pm$ 4.62	20	10.48 $\pm$ 3.68	58	9.80 $\pm$ 3.57	76	10.44 $\pm$ 3.77
Day 30	49	10.20 $\pm$ 6.77	20	11.38 $\pm$ 4.09	20	11.83 $\pm$ 4.32	58	9.93 $\pm$ 3.88	71	10.86 $\pm$ 3.75
Day 90	47	11.64 $\pm$ 7.98	20	11.97 $\pm$ 5.03	20	10.97 $\pm$ 3.18	55	9.56 $\pm$ 3.12	65	11.99 $\pm$ 9.36
Day 120	46	11.71 $\pm$ 7.85	19	12.12 $\pm$ 5.25	20	11.61 $\pm$ 2.58	48	9.63 $\pm$ 3.58	45	11.63 $\pm$ 4.72
Day 180	45	11.12 $\pm$ 7.58	19	11.67 $\pm$ 5.35	19	11.22 $\pm$ 3.44	51	9.19 $\pm$ 2.42	46	11.47 $\pm$ 3.77

### (3) Osteocalcin

Serum osteocalcin was measured by an IRMA from Immutopics (San Clemente, CA). The LLQ was 0.45  $\mu$ g/L. The intra- and inter- assay coefficients were 5.6 and 4.4%, respectively. The UCLA-Harbor Medical Center reports that the normal male adult range for the osteocalcin assay ranges from 2.9 to 12.7  $\mu$ g/L.

As shown in FIG. 17 and Table 20, the baseline mean serum osteocalcin levels were within the normal range in all treatment groups. During the first 90-day treatment, mean serum osteocalcin increased with testosterone replacement in all subjects as a group without significant differences between the groups. With continued treatment serum osteocalcin either plateaued or showed a decrease by day 180.

**Table 20: Osteocalcin Concentrations on Each of the Observation Days  
by Final Treatment Group (Mean  $\pm$  SD)**

	N	5 g/day T-gel	N	5 $\Rightarrow$ 7.5 g/day T-gel	N	10 $\Rightarrow$ 7.5 g/day T-gel	N	10 g/day T-gel	N	T-Patch
Day 0	53	4.62 $\pm$ 1.55	20	5.01 $\pm$ 2.03	20	4.30 $\pm$ 1.28	58	4.58 $\pm$ 1.92	76	4.53 $\pm$ 1.54
Day 30	49	4.63 $\pm$ 1.65	20	5.35 $\pm$ 2.06	20	4.48 $\pm$ 1.72	58	4.91 $\pm$ 2.08	72	5.17 $\pm$ 1.61
Day 90	47	4.91 $\pm$ 2.15	20	5.29 $\pm$ 1.87	19	4.76 $\pm$ 1.50	55	4.83 $\pm$ 2.13	66	5.18 $\pm$ 1.53
Day 120	46	4.95 $\pm$ 1.97	18	4.97 $\pm$ 1.60	20	4.71 $\pm$ 1.39	49	4.61 $\pm$ 2.01	47	4.98 $\pm$ 1.87
Day 180	45	4.79 $\pm$ 1.82	19	4.89 $\pm$ 1.54	19	4.47 $\pm$ 1.49	51	3.76 $\pm$ 1.60	46	5.15 $\pm$ 2.18

### (4) Type I Procollagen

Serum type I procollagen was measured using a RIA kit from Incstar Corp (Stillwater, MN). The LLQ of the procollagen assay was 5  $\mu$ g/L, and the intra- and inter-assay precisions

were 6.6 and 3.6%, respectively. The UCLA-Harbor Medical Center reports that the normal adult male concentration of type I procollagen ranges from 56 to 310 µg/L.

FIG. 18 and Table 21 show that serum procollagen generally followed the same pattern as serum osteocalcin. At baseline the mean levels were similar and within the normal range in all treatment groups. With transdermal treatment, serum procollagen increased significantly in all subjects as a group without treatment group differences. The increase in procollagen was highest on day 30 and then plateaued until day 120. By day 180, the serum procollagen levels returned to baseline levels.

**Table 21: Procollagen Concentrations on Each of the Observation Days  
by Final Treatment Group (Mean ± SD)**

	N	5 g/day T-gel	N	5 => 7.5 g/day T-gel	N	10 => 7.5 g/day T-gel	N	10 g/day T-gel	N	T-Patch
Day 0	53	115.94 ± 43.68	20	109.27 ± 32.70	20	120.93 ± 28.16	58	125.33 ± 57.57	76	122.08 ± 51.74
Day 30	49	141.09 ± 64.02	20	141.41 ± 77.35	20	147.25 ± 49.85	58	149.37 ± 60.61	71	139.26 ± 59.12
Day 90	47	137.68 ± 68.51	20	129.02 ± 60.20	29	144.60 ± 58.20	55	135.59 ± 51.54	66	130.87 ± 49.91
Day 120	46	140.07 ± 81.48	19	133.61 ± 54.09	20	139.00 ± 64.96	50	128.48 ± 45.56	46	130.39 ± 42.22
Day 180	45	119.78 ± 49.02	19	108.78 ± 35.29	19	123.51 ± 39.30	51	108.52 ± 38.98	45	120.74 ± 56.10

**c. Urine Bone Turnover Markers: N-telopeptide/Cr and Ca/Cr Ratios**

Urine calcium and creatinine were estimated using standard clinical chemistry procedures by an autoanalyzer operated by the UCLA-Harbor Pathology Laboratory. The procedures were performed using the COBAS MIRA automated chemistry analyzer system manufactured by Roche Diagnostics Systems. The sensitivity of the assay for creatinine was 8.9 mg/dL and the LLQ was 8.9 mg/dL. According to the UCLA-Harbor Medical Center, creatinine levels in normal adult men range from 2.1 mM to 45.1 mM. The sensitivity of the assay for calcium was 0.7 mg/dL and the LLQ was 0.7 mg/dL. The normal range for urine calcium is 0.21 mM to 7.91 mM.

N-telopeptides were measured by an enzyme-linked immunosorbant assay ("ELISA") from Ostex (Seattle, WA). The LLQ for the N-telopeptide assay was 5 nM bone collagen

equivalent ("BCE"). The intra- and inter-assay had a precision of 4.6 and 8.9%, respectively. The normal range for the N-telopeptide assay was 48-2529 nM BCE. Samples containing low or high serum/urine bone marker levels were reassayed after adjusting sample volume or dilution to ensure all samples would be assayed within acceptable precision and accuracy.

5 The normal adult male range for the N-telopeptide/Cr ratio is 13 to 119 nM BCE/nM Cr. As shown in FIG. 19 and Table 22, urinary N-telopeptide/Cr ratios were similar in all three treatment groups at baseline but decreased significantly in the AndroGel® 10.0 g/day group but not in the AndroGel® 5.0 g/day or testosterone patch group during the first 90 days of treatment. The decrease was maintained such that urinary N-telopeptide/Cr ratio remained lower than baseline in AndroGel® 10.0 g/day and in those subjects adjusted to 7.5 g/day from 10.0 g/day group at day 180. This ratio also decreased in the testosterone patch treatment group by day 180.

**Table 22: N-Telopeptide/Cr Ratio on Each of the Observation Days by Initial Treatment Group (Mean ± SD)**

Initial Treatment Group	N	5.0 g/day T-gel	N	10.0 g/day T-gel	N	T-Patch	Across-group p-value
Day 0	71	90.3 ± 170.3	75	98.0 ± 128.2	75	78.5 ± 82.5	0.6986
Day 30	65	74.6 ± 79.3	73	58.4 ± 66.4	66	91.6 ± 183.6	0.3273
Day 90	62	70.4 ± 92.6	73	55.2 ± 49.1	63	75.0 ± 113.5	0.5348
Day 120	35	78.8 ± 88.2	36	46.6 ± 36.4	21	71.2 ± 108.8	0.2866
Day 180	64	68.2 ± 81.1	70	46.9 ± 43.1	47	49.4 ± 40.8	0.2285

15 The normal range for Ca/Cr ratio is 0.022 to 0.745 mM/mM. FIG. 20 shows no significant difference in baseline urinary Ca/Cr ratios in the three groups. With transdermal testosterone replacement therapy, urinary Ca/Cr ratios did not show a significant decrease in any treatment group at day 90. With continued testosterone replacement to day 180, urinary Ca/Cr showed marked variation without significant changes in any treatment groups.

**Table 23: Ca/Cr Ratio on Each of the Observation Days by Initial Treatment Group (Mean ± SD)**

Initial Treatment Group	N	5.0 g/day T-gel	N	10.0 g/day T-gel	N	T-Patch	Across-group p-value
Day 0	71	0.150 ± 0.113	75	0.174 ± 0.222	75	0.158 ± 0.137	0.6925
Day 30	65	0.153 ± 0.182	73	0.128 ± 0.104	66	0.152 ± 0.098	0.3384
Day 90	63	0.136 ± 0.122	73	0.113 ± 0.075	63	0.146 ± 0.099	0.2531

Initial Treatment Group	N	5.0 g.day T-gel	N	10.0 g/day T-gel	N	T-Patch	Across-group p-value
Day 120	36	0.108 ± 0.073	36	0.117 ± 0.090	21	0.220 ± 0.194	0.0518
Day 180	64	0.114 ± 0.088	70	0.144 ± 0.113	47	0.173 ± 0.108	0.0398

Interestingly, the change in Ca/Cr ratio from baseline at day 90 was inversely related to the baseline Ca/Cr ratios. Similarly, the change in urine N-telopeptide/Cr ratio was also inversely proportional to the baseline N-telopeptide/Cr ratio ( $r=-0.80$ ,  $p=0.0001$ ). Thus subjects with the highest bone resorption markers at baseline showed the largest decreases of these markers during transdermal testosterone replacement. The decreases in urinary bone resorption markers were most prominent in subjects who had highest baseline values, suggesting that hypogonadal subjects with the most severe metabolic bone disease responded most to testosterone replacement therapy.

#### d. Serum Calcium

Serum calcium showed no significant inter-group differences at baseline, nor significant changes after testosterone replacement. Serum calcium levels showed insignificant changes during testosterone replacement.

### 3. Libido, Sexual Performance, and Mood

Sexual function and mood were assessed by questionnaires the patients answered daily for seven consecutive days before clinic visits on day 0 and on days 30, 60, 90, 120, 150, and 180 days during gel and patch application. The subjects recorded whether they had sexual day dreams, anticipation of sex, flirting, sexual interaction (*e.g.*, sexual motivation parameters) and orgasm, erection, masturbation, ejaculation, intercourse (*e.g.*, sexual performance parameters) on each of the seven days. The value was recorded as 0 (none) or 1 (any) for analyses and the number of days the subjects noted a parameter was summed for the seven-day period. The average of the four sexual motivation parameters was taken as the sexual motivation score and that of the five sexual motivation parameters as the sexual motivation mean score (0 to 7). The

subjects also assessed their level of sexual desire, sexual enjoyment, and satisfaction of erection using a seven-point Likert-type scale (0 to 7) and the percent of full erection from 0 to 100%. The subjects rated their mood using a 0 to 7 score. The parameters assessed included positive mood responses: alert, friendly, full of energy, well/good feelings and negative mood responses: angry, irritable, sad, tired, nervous. Weekly average scores were calculated. The details of this questionnaire had been described previously and are fully incorporated by reference. See Wang et al., *Testosterone Replacement Therapy Improves Mood in Hypogonadal Men – A Clinical Research Center Study*, 81 J. CLINICAL ENDOCRINOLOGY & METABOLISM 3578-3583 (1996).

**a. Libido**

As shown in FIG. 21(a), at baseline, sexual motivation was the same in all treatment groups. After transdermal testosterone treatment, overall sexual motivation showed significant improvement. The change in the summary score from baseline, however, was not different among the three treatment groups.

Libido was assessed from responses on a linear scale of: (1) overall sexual desire, (2) enjoyment of sexual activity without a partner, and (3) enjoyment of sexual activity with a partner. As shown in FIG. 21(b) and Table 24, as a group, overall sexual desire increased after transdermal testosterone treatment without inter-group difference. Sexual enjoyment with and without a partner (FIG. 21(c) and Tables 25 and 26) also increased as a group.

Similarly the sexual performance score improved significantly in all subjects as a groups. The improvement in sexual performance from baseline values was not different between transdermal preparations.

**Table 24: Overall Sexual Desire  
Changes From Day 0 to Day 180  
by Initial Treatment Group (Mean  $\pm$  SD)**

Initial Treatment Group	N	Day 0	N	Day 180	N	Change From Day 0 to Day 180	Within-Group p-value
5.0 g/day T-gel	69	2.1 $\pm$ 1.6	63	3.5 $\pm$ 1.6	60	1.4 $\pm$ 1.9	0.0001
10.0 g/day T-gel	77	2.0 $\pm$ 1.4	68	3.6 $\pm$ 1.6	67	1.5 $\pm$ 1.9	0.0001
T-Patch	72	2.0 $\pm$ 1.6	47	3.1 $\pm$ 1.9	45	1.6 $\pm$ 2.1	0.0001
Across-Groups p-value		0.8955		0.2247		0.8579	

**Table 25: Level of Sexual Enjoyment Without a Partner  
Changes From Day 0 to Day 180  
by Initial Treatment Group (Mean  $\pm$  SD)**

Initial Treatment Group	N	Day 0	N	Day 180	N	Change From Day 0 to Day 180	Within-Group p-value
5.0 g/day T-gel	60	1.5 $\pm$ 1.9	51	1.9 $\pm$ 1.9	44	0.8 $\pm$ 1.4	0.0051
10.0 g/day T-gel	63	1.2 $\pm$ 1.4	53	2.2 $\pm$ 1.9	48	1.1 $\pm$ 1.6	0.0001
T-Patch	66	1.4 $\pm$ 1.8	44	2.2 $\pm$ 2.3	40	1.0 $\pm$ 1.9	0.0026
Across-Groups p-value		0.6506		0.7461		0.6126	

**Table 26: Level of Sexual Enjoyment With a Partner  
Change from Day 0 to Day 180  
by Initial Treatment Group (Mean  $\pm$  SD)**

Initial Treatment Group	N	Day 0	N	Day 180	N	Change From Day 0 to Day 180	Within-Group p-value
5.0 g/day T-gel	64	2.1 $\pm$ 2.1	55	2.6 $\pm$ 2.2	48	0.4 $\pm$ 2.2	0.0148
10.0 g/day T-gel	66	1.8 $\pm$ 1.7	58	3.0 $\pm$ 2.2	52	1.0 $\pm$ 2.3	0.0053
T-Patch	61	1.5 $\pm$ 1.7	40	2.2 $\pm$ 2.4	35	0.7 $\pm$ 2.3	0.1170
Across-Groups p-value		0.2914		0.1738		0.3911	

#### **b. Sexual Performance**

FIG. 22(a) shows that while all treatment groups had the same baseline sexual performance rating, the rating improved with transdermal testosterone treatment in all groups. In addition, as a group, the subjects' self-assessment of satisfaction of erection (FIG. 22(b) and Table 27) and percent full erection (FIG. 22(c) and Table 28) were also increased with testosterone replacement without significant differences between groups.



The improvement in sexual function was not related to the dose or the delivery method of testosterone. Nor was the improvement related to the serum testosterone levels achieved by the various testosterone preparations. The data suggest that once a threshold (serum testosterone level probably at the low normal range) is achieved, normalization of sexual function occurs.

5 Increasing serum testosterone levels higher to the upper normal range does not further improve sexual motivation or performance.

**Table 27: Satisfaction with Duration of Erection  
Change from Day 0 to Day 180  
by Initial Treatment Group (Mean  $\pm$  SD)**

Initial Treatment Group	N	Day 0	N	Day 180	N	Change From Day 0 to Day 180	Within-Group p-value
5.0 g/day T-gel	55	2.5 $\pm$ 2.1	57	4.3 $\pm$ 1.8	44	1.9 $\pm$ 2.0	0.0001
10/0 g/day T-gel	64	2.9 $\pm$ 1.9	58	4.5 $\pm$ 1.7	53	1.5 $\pm$ 2.0	0.0001
T-Patch	45	3.4 $\pm$ 2.1	34	4.5 $\pm$ 2.0	20	1.3 $\pm$ 2.1	0.0524
Across-Groups p-value		0.1117		0.7093		0.5090	

**Table 28: Percentage of Full Erection  
Change from Day 0 to Day 180  
by Initial Treatment Group (Mean  $\pm$  SD)**

Initial Treatment Group	N	Day 0	N	Day 180	N	Change From Day 0 to Day 180	Within-Group p-value
5.0 g/day T-gel	53	53.1 $\pm$ 24.1	57	67.4 $\pm$ 22.5	43	18.7 $\pm$ 22.1	0.0001
10.0 g/day T-gel	62	59.6 $\pm$ 22.1	59	72.0 $\pm$ 20.2	52	10.4 $\pm$ 23.4	0.0001
T-Patch	47	56.5 $\pm$ 24.7	33	66.7 $\pm$ 26.7	19	12.7 $\pm$ 20.3	0.0064
Across-Groups p-value		0.3360		0.4360		0.1947	

### c. Mood

The positive and negative mood summary responses to testosterone replacement therapy are shown in FIGS. 23(a) and 23(b). All three treatment groups had similar scores at baseline and all subjects as a group showed improvement in positive mood. Similarly, the negative mood summary scores were similar in the three groups at baseline and as a group the responses to transdermal testosterone applications showed significant decreases without showing between group differences. Specifically, positive mood parameters, such as sense of well being and

energy level, improved and negative mood parameters, such as sadness and irritability, decreased. The improvement in mood was observed at day 30 and was maintained with continued treatment. The improvement in mood parameters was not dependent on the magnitude of increase in the serum testosterone levels. Once the serum testosterone increased into the low normal range, maximal improvement in mood parameters occurred. Thus, the responsiveness in sexual function and mood in hypogonadal men in response to testosterone therapy appeared to be dependent on reaching a threshold of serum testosterone at the low normal range.

#### 4. Muscle Strength

Muscle strength was assessed on days 0, 90, and 180. The one-repetitive maximum ("1-RM") technique was used to measure muscle mass in bench press and seated leg press exercises. The muscle groups tested included those in the hips, legs, shoulders, arms, and chest. The 1-RM technique assesses the maximal force generating capacity of the muscles used to perform the test. After a 5-10 minute walking and stretching period, the test began with a weight believe likely to represent the patient's maximum strength. The test was repeated using increments of about 2-10 pounds until the patient was unable to lift additional weight with acceptable form. Muscle strength was assessed in 167 out of the 227 patients. Four out of 16 centers did not participate in the muscle strength testing because of lack of the required equipment.

The responses of muscle strength testing by the arm/chest and leg press tests are shown in FIG. 24(a) and 24(b) and Table 29. There were no statistical significant differences in arm/chest or leg muscle strength among the three groups at baseline. In general, muscle strength improved in both the arms and legs in all three treatment groups without inter-group differences at both day 90 and 180. The results showed an improvement in muscle strength at 90 and 180 days, more in the legs than the arms, which was not different across treatment groups nor on the different days

of assessment. Adjustment of the dose at day 90 did not significantly affect the muscle strength responses to transdermal testosterone preparations.

**Table 29: Muscle Strength – Days 0, 90, and 180 Levels and Change (lbs.)  
from Day 0 to Day 90 and from Day 0 to Day 180  
by Final Treatment Group**

Final Treatment Group	Study Day	Seated Leg Press		Arm/Chest (Bench Press)	
		N	Mean $\pm$ SD (lbs.)	N	Mean $\pm$ SD (lbs.)
5.0 g/day T-gel	0	37	356.8 $\pm$ 170.0	37	100.5 $\pm$ 37.4
	90	30	396.4 $\pm$ 194.3	31	101.2 $\pm$ 30.7
	□ 0-90	30	25.8 $\pm$ 49.2	31	4.0 $\pm$ 10.0
	180	31	393.4 $\pm$ 196.6	31	99.7 $\pm$ 31.4
	□ 0-180	31	19.9 $\pm$ 62.4	31	1.3 $\pm$ 13.0
7.5 g/day T-gel (from 5.0 g/day)	0	16	302.8 $\pm$ 206.5	16	102.8 $\pm$ 48.9
	90	15	299.8 $\pm$ 193.9	15	109.5 $\pm$ 47.6
	□ 0-90	15	17.0 $\pm$ 88.4	15	5.0 $\pm$ 21.3
	180	14	300.6 $\pm$ 203.0	14	108.5 $\pm$ 49.3
	□ 0-180	14	-0.1 $\pm$ 110.2	14	5.6 $\pm$ 30.4
7.5 g/day T-gel (From 10.0 g/day)	0	14	363.4 $\pm$ 173.8	14	123.3 $\pm$ 54.7
	90	14	401.6 $\pm$ 176.6	14	134.6 $\pm$ 57.5
	□ 0-90	14	38.2 $\pm$ 42.9	14	11.3 $\pm$ 10.5
	180	12	409.9 $\pm$ 180.2	14	132.3 $\pm$ 61.5
	□ 0-180	12	33.9 $\pm$ 67.3	14	9.0 $\pm$ 18.7
10.0 g/day T-gel	0	45	345.9 $\pm$ 186.9	43	114.7 $\pm$ 55.1
	90	43	373.5 $\pm$ 194.8	41	119.8 $\pm$ 54.2
	□ 0-90	43	27.6 $\pm$ 45.1	41	4.6 $\pm$ 12.8
	180	36	364.4 $\pm$ 189.1	34	112.0 $\pm$ 45.5
	□ 0-180	36	32.2 $\pm$ 72.3	34	1.9 $\pm$ 14.8
T-Patch	0	55	310.4 $\pm$ 169.7	54	99.2 $\pm$ 43.1
	90	46	344.9 $\pm$ 183.9	46	106.2 $\pm$ 44.0
	□ 0-90	46	25.4 $\pm$ 37.0	46	3.2 $\pm$ 12.0
	180	36	324.8 $\pm$ 199.0	35	104.8 $\pm$ 44.8
	□ 0-180	36	15.2 $\pm$ 54.7	35	2.3 $\pm$ 15.7

## 5. Body Composition

Body composition was measured by DEXA with Hologic 2000 or 4500A series on days 0, 90, and 180. These assessments were done in 168 out of 227 subjects because the Hologic DEXA equipment was not available at 3 out of 16 study centers. All body composition measurements were centrally analyzed and processed by Hologic (Waltham, MA).

At baseline, there were no significant differences in total body mass ("TBM"), total body lean mass ("TLN"), percent fat ("PFT"), and total body fat mass ("TFT") in the three treatment groups. As shown in FIGS. 25(a) and Table 30, all treatment groups incurred an overall increase in TBM. The increase in TBM was mainly due to the increases in TLN. FIG. 25(b) and Table 30 show that after 90 days of testosterone replacement the increase in TLN was significantly higher in the 10.0 g/day AndroGel<sup>®</sup> group than in the other two groups. At day 180, the increases in TLN were further enhanced or maintained in all AndroGel<sup>®</sup> treated groups, as well as in the testosterone patch group.

FIGS. 25(c) and (d) show that the TFT and the PFT decreased in all transdermal AndroGel<sup>®</sup> treatment groups. At 90 days of treatment, TFT was significantly reduced by in the 5.0 g/day and 10.0 g/day AndroGel<sup>®</sup> groups, but was not changed in the testosterone patch group. This decrease was maintained at day 180. Correspondingly, at the 90 and 180, the decrease in PFT remained significantly lower in all AndroGel<sup>®</sup> treated groups but not significantly reduced in the testosterone patch group.

The increase in TLN and the decrease in TFT associated with testosterone replacement therapy showed significant correlations with the serum testosterone level attained by the testosterone patch and the different doses of AndroGel<sup>®</sup>. Testosterone gel administered at 10.0 g/day increased lean mass more than the testosterone patch and the 5.0 g/day AndroGel<sup>®</sup> groups. The changes were apparent on day 90 after treatment and were maintained or enhanced at day 180. Such changes in body composition was significant even though the subjects were withdrawn from prior testosterone therapy for six weeks. The decrease in TFT and PFT was also related to the serum testosterone achieved and were different across the treatment groups. The testosterone patch group did not show a decrease in PFT or TFT after 180 days of treatment.

Treatment with AndroGel® (50 to 10.0 g/day) for 90 days reduced PFT and TFT. This decrease was maintained in the 5.0 and 7.5 g/day groups at 180 days but were further lowered with continued treatment with the higher dose of the AndroGel®.

**Table 30: Mean Change in Body Composition Parameters (DEXA)  
From Baseline to Day 90 and Baseline to Day 180  
By Final Treatment Groups**

Final Treatment Group	Mean Change from Day 0 – Day 90				
	N	TFT (g)	TLN (g)	TBM (g)	PFT
5.0 g/day T-gel	43	-782 ± 2105	1218 ± 2114	447 ± 1971	-1.0 ± 2.2
7.5 g/day (from 5.0 g/day)	12	-1342 ± 3212	1562 ± 3321	241 ± 3545	-1.0 ± 3.1
7.5 g/day (from 10.0 g/day)	16	-1183 ± 1323	3359 ± 2425	2176 ± 2213	-2.0 ± 1.5
10.0 g/day T-gel	45	-999 ± 1849	2517 ± 2042	1519 ± 2320	-1.7 ± 1.8
T-Patch	52	11 ± 1769	1205 ± 1913	1222 ± 2290	-0.4 ± 1.6

Final Treatment Group	Mean Change from Day 0 – Day 180				
	N	TFT (g)	TLN (g)	TBM (g)	PFT
5.0 g/day T-gel	38	-972 ± 3191	1670 ± 2469	725 ± 2357	-1.3 ± 3.1
7.5 g/day (from 5.0 g/day)	13	-1467 ± 3851	2761 ± 3513	1303 ± 3202	-1.5 ± 3.9
7.5 g/day (from 10.0 g/day)	16	-1333 ± 1954	3503 ± 1726	2167 ± 1997	-2.2 ± 1.7
10.0 g/day T-gel	42	-2293 ± 2509	3048 ± 2284	771 ± 3141	-2.9 ± 2.1
T-Patch	34	293 ± 2695	997 ± 2224	1294 ± 2764	-0.3 ± 2.2

## 6. Lipid profile and blood chemistry

The serum total, HDL, and LDL cholesterol levels at baseline were not significantly different in all treatment groups. With transdermal testosterone replacement, there were no overall treatment effects nor inter-group differences in serum concentrations of total, HDL- and LDL-cholesterol (FIG. 5(d)) and triglycerides (data not shown). There was a significant change of serum total cholesterol concentrations as a group with time ( $p=0.0001$ ), the concentrations on day 30, 90, and 180 were significantly lower than day 0.

Approximately 70 to 95% of the subjects had no significant change in their serum lipid profile during testosterone replacement therapy. Total cholesterol levels which were initially

high were lowered into the normal range (of each center's laboratory) at day 180 in 17.2, 20.4, and 12.2% of subjects on testosterone patch, AndroGel® 5.0 g/day and AndroGel® 10.0 g/day, respectively. Serum HDL-cholesterol levels (initially normal) were reduced to below the normal range (of each center's laboratory) in 9.8, 4.0, 9.1, and 12.5% of subjects at day 180 in the testosterone patch, AndroGel® 5.0, 7.5, and 10.0 g/day groups, respectively. There was no clinically significant changes in renal or liver function tests in any treatment group.

## 7. Skin Irritations

Skin irritation assessments were performed at every clinic visit using the following scale:

0 = no erythema; 1 = minimal erythema; 2 = moderate erythema with sharply defined borders; 3 = intense erythema edema; and 4 = intense erythema with edema and blistering/erosion.

Tolerability of the daily application of AndroGel® at the tested dosages was much better than with the permeation-enhanced testosterone patch. Minimal skin irritation (erythema) at the application site was noted in three patients in the AndroGel® 5.0 g/day group (5.7%) and another three in the AndroGel® 10.0 g/day group (5.3%). Skin irritation varying in intensity from minimal to severe (mild erythema to intense edema with blisters) occurred in 65.8% of patients in the patch group. Because of the skin irritation with the testosterone patch, 16 subjects discontinued the study; 14 of these had moderate to severe skin reactions at the medication sites. No patients who received AndroGel® discontinued the study because of adverse skin reactions. The open system and the lower concentration of alcohol in the AndroGel® formulation markedly reduced skin irritation resulting in better tolerability and continuation rate on testosterone replacement therapy.

Moreover, based on the difference in the weight of the dispensed and returned AndroGel® bottles, the mean compliance was 93.1% and 96.0% for the 5.0 g/day and 10.0 g/day AndroGel® groups during days 1-90, respectively. Compliance remained at over 93% for the three

AndroGel® groups from days 91-180. In contrast, based on counting the patches returned by the subjects, the testosterone patch compliance was 65% during days 1-90 and 74% during days 91-180. The lower compliance in the testosterone patch group was mostly due to skin reactions from the subjects' records.

**Table 31: Incidence of Skin-Associated Adverse Events: Day 1 to Day 180 in Patients Who Remained on Initial Treatment**

	5.0 g/day T-gel N = 53	10.0 g/day T-gel N = 57	T-Patch N = 73
Total	16 (30.2%)	18 (31.6%)	50 (68.5%)
Application Site Reaction	3 (5.7%)	3 (5.3%)	48 (65.8%)
Acne	1 (1.9%)	7 (12.3%)	3 (4.1%)
Rash	4 (7.5%)	4 (7.0%)	2 (2.7%)
Skin Disorder	2 (3.8%)	1 (1.8%)	1 (1.4%)
Skin Dry	2 (3.8)	0 (0.0%)	1 (1.4%)
Sweat	0 (0.0%)	2 (3.5%)	0 (0.0%)
Reaction Unevaluable	2 (3.6%)	1 (1.7%)	0 (0.0%)
Cyst	0 (0.0%)	0 (0.0%)	2 (2.7%)

#### Example 2: Gel Delivery Dosage Forms and Devices

The present invention is also directed to a method for dispensing and packaging the gel. In one embodiment, the invention comprises a hand-held pump capable of delivering about 2.5 g of testosterone gel with each actuation. In another embodiment, the gel is packaged in foil packets comprising a polyethylene liner. Each packet holds about 2.5 g of testosterone gel. The patient simply tears the packet along a perforated edge to remove the gel. However, because isopropyl myristate binds to the polyethylene liner, additional isopropyl myristate is added to the gel in order to obtain a pharmaceutically effective gel when using this delivery embodiment. Specifically, when dispensing the gel via the foil packet, about 41% more isopropyl myristate is used in the gel composition (*i.e.*, about 0.705 g instead of about 0.5 g in Table 5), to compensate for this phenomenon.

The composition can also be dispensed from a rigid multi-dose container (e.g., with a hand pump) having a larger foil packet of the composition inside the container. Such larger packets also comprise a polyethylene liner as above.

Both embodiments permit a patient to deliver accurate but incremental amounts of gel (e.g., either 2.5 g, 5.0 g, 7.5 g, etc.) to the body. These delivery mechanisms thus permit the gel to be administered in unit dose form depending on the particular needs and characteristics of the patient.

Although the invention has been described with respect to specific embodiments and examples, it should be appreciated that other embodiments utilizing the concept of the present invention are possible without departing from the scope of the invention. The present invention is defined by the claimed elements, and any and all modifications, variations, or equivalents that fall within the true spirit and scope of the underlying principles.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
220